



Appendix A

Sea Level Rise Science and Projections for Future Change

DRIVERS OF SEA LEVEL RISE

The main mechanisms driving increases in *global* sea level are: 1) expansion of sea water as it gets warmer (thermal expansion) and, 2) increases in the amount of water in the ocean from melting of land-based glaciers and ice sheets as well as human-induced changes in water storage and groundwater pumping (Chao *et al.* 2008; Wada *et al.* 2010; Konikow 2011).¹ The reverse processes can cause global sea level to fall.

Sea level at the *regional and local levels* often differs from the average global sea level.² Regional variability in sea level results from large-scale tectonics and ocean and atmospheric circulation patterns. The primary factors influencing local sea level include tides, waves, atmospheric pressure, winds, vertical land motion and short duration changes from seismic events, storms, and tsunamis. Other determinants of local sea level include changes in the ocean floor (Smith and Sandwell 1997), confluence of fresh and saltwater, and proximity to major ice sheets (Clark *et al.* 1978; Perette *et al.* 2013).

Over the long-term, sea level trends in California have generally followed global trends (Cayan *et al.* 2009; Cayan *et al.* 2012). However, global projections do not account for California's regional water levels or land level changes. California's water levels are influenced by large-scale oceanographic phenomena such as the El Niño Southern Oscillation (ENSO) and the Pacific Decadal Oscillation (PDO), which can increase or decrease coastal water levels for extended periods of time. [Figure A-1](#) shows how El Niño and La Niña events have corresponded to mean sea level in California in the past. California's land levels are also affected by plate tectonics and earthquakes. Changes to water as well as land levels are important factors in regionally down-scaled projections of future sea level. It follows that the sea level rise projections specific to California are more relevant to efforts in the coastal zone of California than projections of global mean sea level.

¹ Large movements of the tectonic plates have been a third major mechanism for changes in global sea level. The time periods for plate movements to significantly influence global sea level are beyond the time horizons used for even the most far-reaching land-use decisions. Plate dynamics will not be included in these discussions of changes to future sea level.

² For further discussion of regional sea level variations and regional sea level rise projections, see Yin *et al.* 2010, Slangen *et al.* 2012, and Levermann *et al.* 2013, as examples.




Figure A-1. Variations in monthly mean sea level at Fort Point, San Francisco, 1854 to 2013. Mean sea level heights (in ft) are relative to mean lower low water (MLLW). Purple line represents the 5-year running average. Note that the monthly mean sea level has varied greatly throughout the years and that several of the peaks occurred during strong El Niño events (red highlight). Periods of low sea level often occurred during strong La Niña events (blue highlight). The current “flat” sea level condition can also be seen in the 5-year running average. (Sources: NOAA CO-OPS data, Station 9414290, <http://tidesandcurrents.noaa.gov/> (sea level); NOAA Climate Prediction Center, <http://www.elnino.noaa.gov/> (ENSO data))

APPROACHES FOR PROJECTING FUTURE GLOBAL SEA LEVEL RISE

This section provides an overview of some of the more well-known approaches that have been used to project sea level changes and their relevance to California. [Appendix B](#) will cover how these projections can be used to determine water conditions at the local scale.

There is no single, well-accepted technique for projecting future sea level rise. Understanding future sea level rise involves projecting future changes in glaciers, ice sheets, and ice caps, as well as future groundwater and reservoir storage. Two subjects in particular present challenges in sea level rise modeling. First, future changes to glaciers, ice sheets, and ice caps are not well understood and, due to the potential for non-linear responses from climate change, they present many difficulties for climate models (Overpeck 2006; Pfeffer *et al.* 2008; van den Broecke *et al.* 2011; Alley and Joughin 2012; Shepherd *et al.* 2012; Little *et al.* 2013). Second, the actual magnitudes of the two human-induced changes – pumping of groundwater and storage of water in reservoirs – are poorly quantified, but the effects of these activities are understood and can be modeled (Wada *et al.* 2010). Despite these challenges, sea level rise projections are needed for many coastal management efforts and scientists have employed a variety of techniques to model sea level rise, including:

1. Extrapolation of historical trends;
2. Modeling the physical conditions that cause changes in sea level;
3. Empirical or semi-empirical methods; and
4. Expert elicitations

There are strengths and weaknesses to each approach, and users of any sea level rise projections should recognize that there is no perfect approach for anticipating future conditions. This section provides users of the Guidance document with a general understanding of several of the most widely used sea level rise projection methodologies and their respective advantages and disadvantages. [Figure A-2](#) provides a visual summary of several of the more commonly cited projections of future global and regional sea level rise.




Figure A-2. Sea level rise projections for year 2100 from scientific literature. Graphic summary of the range of average sea level rise (SLR) projections by end of century (2090–2100) from the peer-reviewed literature as compared to the recent National Research Council report for California, Oregon and Washington. The light blue shaded boxes indicate projections for California. Ranges are based on the IPCC scenarios, with the low range represented by the B1 scenario (moderate growth and reliance in the future on technological innovation and low use of fossil fuels) and the high part of the range represented by the A1FI scenario (high growth and reliance in the future on fossil fuels). Details on the methods used and assumptions are provided in the original references.

Extrapolation of Historical Trends

Extrapolation of historical trends in sea level has been used for many years to project future changes in sea level. The approach assumes that there will be no abrupt changes in the processes that drive the long-term trend, and that the driving forces will not change. However, drivers of climate change and sea level rise, such as radiative forcing, are known to be changing, and this method is no longer considered appropriate or viable in climate science.

A recent modification to the historical trend method discussed above has been to estimate rates of sea level rise during the peak of the last interglacial (LIG) period (~125,000 years before present, when some drivers of sea level rise were similar to those today)³ and use these as proxy records to project sea level rise rates to the 21st Century. For example, Katsman *et al.* (2011) and Vellinga *et al.* (2008) used the reconstructed LIG record of sea level change (from Rohling *et al.* 2008) to reconstruct sea level rise rates during rapid climate warming, and applied these rates to estimate sea level at years 2100 and 2200. Similarly, Kopp *et al.* (2009) used sea level rise rates inferred from the LIG to estimate a range of sea level rise for Year 2100 between 1-3 ft (0.3-1 m). Compared to traditional historical trend extrapolation, this modified approach has the advantage of including the dynamic responses of ice sheets and glaciers to past global climates that were significantly warmer than the present, but is limited by the large uncertainties associated with proxy reconstructions of past sea level.

Physical Models

Physical climate models use mathematical equations that integrate the basic laws of physics, thermodynamics, and fluid dynamics with chemical reactions to represent physical processes such as atmospheric circulation, transfers of heat (thermodynamics), development of precipitation patterns, ocean warming, and other aspects of climate. Some models represent only a few processes, such as the dynamics of ice sheets or cloud cover. Other models represent larger scale atmospheric or oceanic circulation, and some of the more complex General Climate Models (GCMs) include atmospheric and oceanic interactions.

Physical models of sea level changes account for the thermal expansion of the ocean and the transfer of water currently stored on land, particularly from glaciers and ice sheets (Church *et al.* 2011). Currently, coupled Atmosphere-Ocean General Circulation Models (AOGCMs) and ice sheet models are replacing energy-balance climate models as the primary techniques supporting sea level projections (IPCC 2013). Ocean density, circulation and sea level are dynamically connected in AOGCMs as critical components of the models include surface wind stress, heat transfer between air and sea, and freshwater fluxes. AOGCM climate simulations have recently been used as input for glacier models (Marzeion *et al.* 2012) which project land-water contributions to sea level.

The Intergovernmental Panel on Climate Change (IPCC) is one of the main sources of peer-reviewed, consensus-based modeling information on climate change. The IPCC does not undertake climate modeling, but uses the outputs from a group of climate models that project

³ During the last interglacial, global mean temperature was 1-2°C warmer than the pre-industrial era (Levermann *et al.* 2013), while global mean sea level was likely 16.4-29.5 ft (5-9 m) above present mean sea level (Kopp *et al.* 2009; Dutton and Lambeck 2012; Levermann *et al.* 2013).

future temperature, precipitation patterns, and sea level rise, based on specific emission scenarios. Early in the 1990s, the IPCC developed basic model input conditions to ensure comparable outputs from the various models. The IPCC initially developed scenarios of future emissions, based on energy development, population and economic growth, and technological innovation. Four families of scenarios (A1, A2, B1, and B2) and subgroups (A1B, A1FI, A1T) were developed and used for climate and sea level rise projections for early IPCC reports (1990, 1995, 2001, 2007). IPCC used 4 new scenarios for the [5th Assessment Report](#) (AR5) in 2013, based on Representative Concentration Pathways (RCPs) that are different greenhouse gas concentration trajectories. These trajectories bear similarities to, but are not directly comparable to the earlier emission scenarios. Projections in IPCC AR5 (2013) differ from the earlier IPCC projections due to improvements in climate science, changes due to the new scenarios, and changes in the models to accommodate the new inputs, with improvements in climate science and model capabilities driving the bulk of the changes.

One finding of the earlier 2007 IPCC report called for improved modeling of ice dynamics. Focused research on ice dynamics to improve the ability of climate models to address the scale and dynamics of change to glaciers, ice sheets, and ice caps was subsequently undertaken (*e.g.*, Price *et al.* 2011; Shepherd *et al.* 2012; Winkelmann *et al.* 2012; Bassis and Jacobs 2013; Little *et al.* 2013). Recent modeling results presented in the AR5 (IPCC 2013) reflect the scientific community's increased understanding in, as well as advances in modeling of the impacts of glacier melting and ocean thermal expansion on sea level change. AR5 scenarios reflect a greater range of global sea level rise (28-98 cm) based on improved modelling of land-ice contributions.

Semi-Empirical Method

The semi-empirical method for projecting sea level rise is based on developing a relationship between sea level and some factor (a proxy) – often atmospheric temperature or radiative forcing – and using this relationship to project changes to sea level. An important aspect for the proxy is that there is fairly high confidence in models of its future changes; a key assumption that is made by this method is that the historical relationship between sea level and the proxy will continue into the future. One of the first projections of this kind was based on the historical relationship between global temperature changes and sea level changes (Rahmstorf 2007). This semi-empirical approach received widespread recognition for its inclusion of sea level rise projections. These projections looked at the temperature projections for two of the previous IPCC (2007) emission scenarios that span the likely future conditions within the report's framework – B1, an optimistic, low-greenhouse gas emission future, and A1FI, a more “business-as-usual” fossil fuel intensive future.⁴ The Rahmstorf 2007 sea level rise projections were used in the California 2009 *Climate Change Scenarios Assessment* (Cayan 2009).

Since the initial semi-empirical projections for future sea level rise (Rahmstorf 2007), other researchers have published different projections based on the IPCC scenarios, using different

⁴ When the IPCC began examining climate change, the available models used a broad range of inputs. In an attempt to evaluate the different model outputs based on the different model characteristics rather than the inputs, the IPCC developed a number of standard greenhouse gas emission scenarios. These scenarios are described in *Response Strategies Working Group III* (IPCC 1990). In general, the B1 scenario projects the lowest temperature and sea level increases and the A1FI projects the highest increases.

data sets or best-fit relationships.⁵ Notably, Vermeer and Rahmstorf (2009) prepared a more detailed methodology that includes both short-term responses and longer-term responses between sea level rise and temperature. These 2009 projections of sea level rise were used in the *Interim Guidance on Sea Level Rise* (OPC 2010) and the California 2012 *Vulnerability and Assessment Report* (Cayan 2012).

There are also several new semi-empirical sea level rise projections based on scenarios other than those developed by the IPCC. For instance, Katsman *et al.* (2011) use a “hybrid” approach that is based on one of the newer radiative forcing scenarios and empirical relationships between temperature change and sea level. Future projections were then modified to include contributions from the melting of major ice sheets based on expert judgment⁶. This yields what they call “high end” SLR projections for Years 2100 and 2200 under several emissions scenarios.

Zecca and Chiari (2012) produced semi-empirical sea level rise projections based on their own scenarios of when fossil fuel resources would be economically exhausted. Though based on a different set of assumptions about human behavior/choices, in terms of global temperature and radiative forcing, the scenarios do not differ greatly from the IPCC scenarios. The results are identified as being “lower bound” sea level rise projections for high, medium, low fuel use scenarios, and “mitigation” (extreme and immediate action to replace fossil fuel use) scenarios. The report then provides projections for the 2000-2200 time period.

Expert Elicitation

Expert elicitation is one of the newer methods that have been used for projecting or narrowing ranges of future sea level rise. Using expert judgment has been an important aspect of scientific inquiry and the scientific method. The method of expert elicitation is a formalized use of experts in climate science and sea level change to help either narrow uncertainty for sea level projections, or to help with specifying extremes of a range. The elicitation method normally begins with experts refining model output information. One of the first attempts to use expert elicitation for sea level rise was a study by Titus and Narayanan (1996), when it was thought there was only 1% probability that sea level would exceed 3.3 ft (1 m) by Year 2100. In 2011, the Arctic Monitoring and Assessment Programme Report (AMAP 2011) surveyed the climate literature to construct a range of estimates of sea level rise by the year 2100, and then used a panel of experts to decide on a smaller, more plausible range. Not surprisingly, the projections supported by the AMAP experts fell right in the middle of the range shown in [Figure A-2](#). Bamber and Aspinall (2013) used a statistical analysis of a large number of expert estimates to

⁵ Semi-empirical projections of sea level rise using relationships between water level and radiative forcing such as those from Grinsted *et al.* (2009), Jevrejeva *et al.* (2010), Katsman *et al.* (2011), Meehl *et al.* (2012), Rahmstorf *et al.* (2012), Schaeffer *et al.* (2012), and Zecca and Chiari (2012) have shown general agreement with the projections by Vermeer and Rahmstorf (2009). The Grinsted *et al.* projections have a wider range than those of Vermeer and Rahmstorf, while the Jevrejeva *et al.* projections are slightly lower. All semi-empirical methods project that sea level in Year 2100 is likely to be much higher than linear projections of historical trends and the projections from the 2007 IPCC.

⁶ Expert judgment has long been part of the scientific process. Expert elicitation, which is a formalized process for using expert judgment, has grown in importance and is discussed as a separate approach for projecting future sea level rise.

develop their projected range of future sea level, projecting sea level rise by 2100 ranging from 1–4.3 ft (0.33–1.32 m), under one of the intermediate AR5 scenarios (RCP 4.5).

Horton *et al.* (2014) surveyed experts in sea level science, based upon published papers, to develop a probabilistic assessment of long-term sea level rise (by the years 2100 and 2300), assuming two very different scenarios. Under one scenario, aggressive efforts would limit greenhouse gas concentrations that would cause global temperature to increase slightly until about 2050 when it would slowly drop (AR5’s RCP 3 scenario). Under the other scenario, temperatures would continue to increase through to 2300 (AR5’s RCP 8.5 scenario). Experts determined that it is likely that sea level rise could remain below 3.3 ft (1 m) for the low emission scenario (RCP 2.6), but that the likely range of future sea level rise for the high emission scenario (RCP 8.5) could be 6.6-9.8 ft (2-3 m).

Kopp *et al.* (2014) have combined detailed process modeling, community assessments and expert elicitation to assign probability distributions of local sea level rise through 2200 for identified communities around the world. Under the high concentration scenario, RCP 8.5, Kopp *et al.* estimate the “maximum physically possible rate of sea level rise” to be 8.2 ft (2.5 m) for the year 2100. This study also finds that sea level rise along the Pacific Coast of the US is close to the global average, and the likely range of sea level is 2-3.3 ft (0.6-1.0 m) by the year 2100 at San Francisco, under the high concentration scenario. In contrast, in areas of high subsidence such as Galveston, Texas, the likely range of sea level in by 2100 ranges from 3.3 to 5 ft (1.0-1.5 m). And, at many of the localities that were examined, including San Francisco, the current 1-in-10 year flooding event is likely to occur every other year by 2100 (five times more frequently) due to sea level rise; the frequency of the 1-in-100 year event is expected to double by the year 2100 with sea level rise.

Coastal communities cannot ignore sea level rise in long-term planning, permitting and project design. The four different approaches to projecting future sea level rise all have varying strengths and weaknesses. As noted earlier in this section, projections, like models, will not be completely accurate, but they are important tools for evaluation nonetheless⁷. The most commonly cited projections provide future sea level as a range, as a way to allow for many of the uncertainties that are part of future climate change. Often, projections of sea level rise rely upon multiple approaches. For example, the 2012 National Research Council (NRC) report was developed through expert judgment that combined information from both physical models and semi-empirical projections.

⁷ George E.P. Box, mathematician and statistician is quoted as saying, “Essentially all models are wrong, but some are useful.”

BEST AVAILABLE SCIENCE ON SEA LEVEL RISE

Global Projections of Sea Level Rise

The best available science on *global* sea level rise projections is currently the IPCC *Fifth Assessment Report: Climate Change 2013* (AR5) released in September 2013. The new report now projects a more rapid sea level rise than the *Fourth Assessment* (AR4) released in 2007. By Year 2100, the AR5 projects global sea level to be more than 50% higher (26-98 cm) than the old projections (18-59 cm) when comparing similar emission scenarios and time periods. The increase in AR5 sea level projections results from improved modelling of land-ice contributions. Substantial progress in the assessment of extreme weather and climate events has also been made since the AR4 as models now better reproduce phenomena like the El Niño-Southern Oscillation (ENSO; IPCC 2013).

National Projections of Sea Level Rise

The [third National Climate Assessment](#) (NCA) was released in May 2014 (Melillo *et al.*), and includes the current best-available science on climate change and sea level rise *at the national scale*.⁸ The sea level rise projections in the NCA were informed by the 2012 NOAA report titled [Global Sea Level Rise Scenarios for the United States National Climate Assessment](#) (Parris *et al.*). This report provides a set of four scenarios of future global sea level rise, as well as a synthesis of the scientific literature on global sea level rise. The NOAA Climate Program Office produced the report in collaboration with twelve contributing authors.⁹ The report includes the following description of the four scenarios of sea level rise by the year 2100:

- **Low scenario:** The lowest sea level change scenario (a rise of 8 in (20 cm)) is based on historical rates of observed sea level change.
- **Intermediate-low scenario:** The intermediate-low scenario (a rise of 1.6 ft (0.5 m)) is based on projected ocean warming.
- **Intermediate- high scenario:** The intermediate-high scenario (a rise of 3.9 ft (1.2 m)) is based on projected ocean warming and recent ice sheet loss.
- **High scenario:** The highest sea level change scenario (a rise of 6.6 ft (2 m)) reflects ocean warming and the maximum plausible contribution of ice sheet loss and glacial melting.

The Parris *et al.* (2012) report recommends that the highest scenario be considered in situations where there is little tolerance for risk. It also provides steps for planners and local officials to modify these scenarios to account for local conditions. These steps are intended for areas where local sea level rise projections have not been developed. For California, the 2018 OPC SLR Guidance report (below) provides scenarios that have been refined for use at the local level, and the Coastal Commission recommends using the OPC projections rather than the global or national scenarios.

⁸ Note that the 4th National Climate Assessment is due to be released in late 2018. <https://www.globalchange.gov/nca4>

⁹ Authors include NOAA, NASA, the US Geologic Survey, the Scripps Institution of Oceanography, the US Department of Defense, the US Army Corps of Engineers, Columbia University, the University of Maryland, the University of Florida, and the South Florida Water Management District.

California-Specific Projections of Sea Level Rise and Best Available Science

The State of California has long-supported the development of scientific information on climate change and sea level rise to help guide planning and decision-making. For example, the State helped support the development of the 2012 National Research Council (NRC) report, [*Sea-Level Rise for the Coasts of California, Oregon, and Washington: Past, Present, and Future*](#), which provided an examination of global and regional sea level rise trends and projections of future sea level. This report was then incorporated into the Ocean Protection Council's 2013 *State Sea-Level Rise Guidance*, and was considered the best available science on sea level rise for California.

More recently, and in response to the release of new scientific studies related to sea level rise, Governor Brown directed the OPC to synthesize recent science on sea level rise and incorporate findings into updates to the State Guidance. In April 2017, a working group of OPC's Science Advisory team (comprised mainly of climate researchers at various academic institutions in California and throughout the country) released a report titled [*Rising Seas in California: An Update on Sea-Level Rise Science*](#). The report highlighted seven key findings:

1. *Scientific understanding of sea level rise is advancing at a rapid pace.* Sea level rise projections have increased substantially over the last few years, particularly for late in the 21st century and under high emissions scenarios, due to our evolving understanding of the dynamics of ice sheet loss. However, there is still significant uncertainty regarding these processes.
2. *The direction of sea level change is clear.* Coastal California is already experiencing the impacts of rising sea levels, and impacts will increase in the future.
3. *The rate of ice loss from the Greenland and Antarctic ice sheets is increasing.* Ice sheet loss will soon overtake thermal expansion of seawater as the primary driver of rising sea levels. Due to a variety of ocean circulation dynamics, ice loss from Antarctica, and particularly West Antarctica, has an outsized impact on California compared to the rest of the world ([Figure A-3](#)). Continued research on this dynamic is critical for accurately projecting future sea level rise along our coast.
4. *New scientific evidence has highlighted the potential for extreme sea level rise.* Recent research (e.g., DeConto and Pollard, 2016; Sweet et al., 2017) has found that, if greenhouse gas emissions are not curtailed, glaciological processes could cross thresholds that lead to rapidly accelerating and effectively irreversible ice loss. The probability of this extreme scenario is currently unknown, but its consideration is important. Significant reductions in greenhouse gas emissions may reduce the likelihood of this extreme scenario, but does not completely eliminate the risk. Importantly, it is difficult to determine if the world is on the track for extreme and irreversible ice loss for some time because the processes that drive extreme ice loss in the later part of the century or beyond are different than those that are driving ice loss now.




Figure A-3. Sea level ‘fingerprints’ resulting from the distribution of ice and water around the Earth and ensuing gravitational and rotational effects. The maps depict the relative response of sea-level to the loss of ice mass from (a) Greenland Ice Sheet (GIS) and (b) West Antarctic Ice Sheet (WAIS). The color bar represents the fractional departure of relative sea level rise from that expected given the ice contribution to global mean sea level. For example, when ice is lost from the Greenland Ice Sheet the relative effect on the US West Coast is 75% of the sea-level rise expected from the water volume added to the ocean. By comparison, when ice is lost from the West Antarctic Ice Sheet the US West Coast experiences 125% of sea-level rise from that expected from the water volume added (from Griggs et al. 2017).

5. *Probabilities of specific sea-level increases can inform decisions.* A probabilistic approach to sea level rise projections, combined with a clear articulation of the implications of uncertainty and the decision support needs of affected stakeholders, is the most appropriate approach for use in a policy setting.

The OPC Scientific Working Group utilized a comprehensive probability approach based on Kopp et al. (2014) that estimates both a comprehensive probability distribution and the likelihood of extreme ‘tail’ outcomes. It is important to note that probabilistic projections do not provide probabilities of occurrence of sea level rise, but rather probabilities that the ensemble of climate models used to estimate contributions of sea level rise (from thermal expansion, ice sheet loss, oceanographic conditions etc.) will predict a certain amount of sea level rise.

Note that the probabilistic projections do not consider the H++ extreme ice loss scenario. The extreme ice loss studies were not included in the inputs to the model ensemble, which means the probability distributions may be an underestimate.¹⁰

6. *Current greenhouse gas emissions policy decisions are shaping our coastal future.* Before 2050, differences in SLR projections under different emissions scenarios are minor. After 2050, SLR projections increasingly depend on the trajectory of greenhouse

¹⁰ The 4th California Climate Assessment developed projections that present a broader range of SLR estimates than the Rising Seas science report and the 2018 OPC SLR Guidance. Both programs’ projections are based on estimates of contributions to SLR from primary sources using different methods, including model projections and expert input. However, the 4th Assessment incorporates the findings from the recent studies regarding the potential for rapid loss of Antarctic ice sheets (which results in the H++ scenario of about 10ft. of SLR by 2100) into its probabilistic projections whereas the OPC reports do NOT include this possibility in the probabilistic projections, as explained above.

gas emissions. If greenhouse gas emissions are not curtailed worldwide, we will see significantly higher rates of sea level rise during the second half of the century.

7. *Waiting for scientific certainty is neither a safe nor prudent option.* Taking action today to assess vulnerabilities and identify and implement adaptation strategies will prevent much greater losses than will occur if action is not taken. Taking a precautionary approach that considers high and extreme scenarios is critical for safeguarding the people and resources of coastal California.

This scientific information was incorporated into OPC's [State Sea-Level Rise Guidance: 2018 Update](#). The OPC Guidance includes projection tables for 12 tide gauges along the California coast for each decade from 2030 to 2150. OPC further recommends utilizing three different projection scenarios to guide planning, permitting, investment, and other decisions based on the type of project, its ability to cope with or adapt to sea level rise, and the consequences to the environment and the project associated with sea level rise. The projection table for the San Francisco tide gauge is provided below ([Table A-1](#)), and tables for other California tide gauges are presented in [Appendix G](#). The 2018 OPC SLR Guidance (along with the foundational Rising Seas science report) is currently considered best available science on sea level rise for the State of California.

The Coastal Commission recommends that the low, medium-high, and extreme risk aversion scenarios from the OPC 2018 Sea-Level Rise Guidance be considered in all relevant local coastal planning and coastal development permitting decisions.

Table A-1. Sea Level Rise Projections for the San Francisco Tide Gauge¹¹ (OPC 2018)

Projected Sea Level Rise (in feet): <i>San Francisco</i>			
	Probabilistic Projections (in feet) (based on Kopp et al. 2014)		H++ Scenario (Sweet et al. 2017)
	Low Risk Aversion	Medium-High Risk Aversion	Extreme Risk Aversion
	<i>Upper limit of "likely range" (~17% probability SLR exceeds...)</i>	<i>1-in-200 chance (0.5% probability SLR exceeds...)</i>	<i>Single scenario (no associated probability)</i>
2030	0.5	0.8	1.0
2040	0.8	1.3	1.8
2050	1.1	1.9	2.7
2060	1.5	2.6	3.9
2070	1.9	3.5	5.2
2080	2.4	4.5	6.6
2090	2.9	5.6	8.3
2100	3.4	6.9	10.2
2110*	3.5	7.3	11.9
2120	4.1	8.6	14.2
2130	4.6	10.0	16.6
2140	5.2	11.4	19.1
2150	5.8	13.0	21.9

**Most of the available climate model experiments do not extend beyond 2100. The resulting reduction in model availability causes a small dip in projections between 2100 and 2110, as well as a shift in uncertainty estimates (see Kopp et al., 2014). Use of 2110 projections should be done with caution and acknowledgement of increased uncertainty around these projections.*

¹¹ Probabilistic projections for the height of sea level rise and the H++ scenario are presented. The H++ projection is a single scenario and does not have an associated likelihood of occurrence. Projections are with respect to a baseline year of 2000 (or more specifically, the average relative sea level over 1991-2009). Table is adapted from the 2018 OPC SLR Guidance to present only the three scenarios OPC recommends evaluating. Additionally, while the OPC tables include low emissions scenarios, only high emissions scenarios, which represent RCP 8.5, are included here because global greenhouse gas emissions are currently tracking along this trajectory. The Coastal Commission will continue to update best available science as necessary, including if emissions trajectories change.

REFERENCES: APPENDIX A

- Alley RB, I Joughin. 2012. Modeling ice-sheet flow. *Science* 336(6081): 551-552. [doi: 10.1126/science.1220530](https://doi.org/10.1126/science.1220530).
- Arctic Monitoring and Assessment Programme (AMAP). 2011. *Snow, Water, Ice and Permafrost in the Arctic (SWIPA): Climate Change and the Cryosphere*. Arctic Monitoring and Assessment Programme, Oslo, Norway, 538 pp. <http://www.amap.no/>.
- Bamber JL, WP Aspinall. 2013. An expert judgment assessment of future sea level rise from the ice sheets. *Nature Climate Change* 3: 424-427. doi:10.1038/nclimate1778.
- Bassis JN, S Jacobs. 2013. Diverse calving patterns linked to glacier geometry. *Nature Geoscience* 6: 833-836. doi: 10.1038/ngeo1887.
- Bedsworth, L, D Cayan, G Franco, L Fisher, S Ziaja. (CA Governor's Office of Planning and Research, Scripps Institution of Oceanography, CA Energy Commission, CA Public Utilities Commission). 2018. Statewide Summary Report. California's Fourth Climate Change Assessment. <http://www.climateassessment.ca.gov/state/index.html>
- California Natural Resources Agency (CNRA). 2009. *California Climate Adaptation Strategy*. http://resources.ca.gov/docs/climate/Statewide_Adaptation_Strategy.pdf.
- Cayan D, M Tyree, M Dettinger, H Hidalgo, T Das, E Maurer, P Bromirski, N Graham, R Flick. 2009. *Climate Change Scenarios and Sea Level Rise Estimates for the California 2009 Climate Change Scenarios Assessment*. California Climate Change Center, CEC-500-2009-014-F. <http://www.energy.ca.gov/2009publications/CEC-500-2009-014/CEC-500-2009-014-F.PDF>.
- Cayan D, M Tyree, D Pierce, T Das. 2012. *Climate Change and Sea Level Rise Scenarios for California Vulnerability and Adaptation Assessment*. A White Paper from the California Energy Commission's California Climate Change Center, CEC-500-2012-008, <http://www.energy.ca.gov/2012publications/CEC-500-2012-008/CEC-500-2012-008.pdf>.
- Chao BF, YH Wu, YS Li. 2008. Impact of artificial reservoir water impoundment on global sea level. *Science* 320(5873): 212. [doi: 10.1126/science.1154580](https://doi.org/10.1126/science.1154580).
- Church JA, NJ White, LF Konikow, CM Domingues, JG Cogley, E Rignot, JM Gregory, MR van der Broeke, AJ Monaghan, I Velicogna. 2011. Revisiting the Earth's sea-level and energy budgets from 1961 to 2008. *Geophysical Research Letters* 40(15): 4066. [doi: 10.1029/2011GL048794](https://doi.org/10.1029/2011GL048794).
- Clark JA, WE Farrell, WR Peltier. 1978. Global changes in postglacial sea level: a numerical calculation. *Quaternary Research* 9(3): 265-287. [doi:10.1016/0033-5894\(78\)90033-9](https://doi.org/10.1016/0033-5894(78)90033-9).
- DeConto RM, Pollard D. 2016. Contribution of Antarctica to past and future sea-level rise. *Nature* 531: 591-7
- Deltacommissie. 2008. *Working together with water: A living land builds for its future*. Findings of the Deltacommissie 2008. http://www.deltacommissie.com/doc/deltareport_summary.pdf.

- Dutton A, K Lambeck. 2012. Ice volume and sea level during the last interglacial. *Science* 337(6091): 216–219. [doi: 10.1126/science.1205749](https://doi.org/10.1126/science.1205749).
- Griggs, G, Árvai, J, Cayan, D, DeConto, R, Fox, J, Fricker, HA, Kopp, RE, Tebaldi, C, Whiteman, EA (California Ocean Protection Council Science Advisory Team Working Group). [Rising Seas in California: An Update on Sea-Level Rise Science](#). California Ocean Science Trust, April 2017.
- Grinsted A, J Moore, S Jevrejeva. 2009. Reconstructing sea level from paleo and projected temperatures 200 to 2100 AD. *Climate Dynamics* 34: 461- 472. [doi:10.1007/s00382-008-0507-2](https://doi.org/10.1007/s00382-008-0507-2).
- Horton BP, S Rahmstorf, SE Engelhart, AC Kemp. 2014. Expert assessment of sea-level rise by AD 2100 and AD 2300. *Quaternary Science Review* 84: 1-6. [doi: 10.1016/j.quascirev.2013.11.002](https://doi.org/10.1016/j.quascirev.2013.11.002).
- Horton R, C Herweijer, C Rosenzweig, J Liu, V Gornitz, AC Ruane. 2008. Sea level rise projections for current generation CGCMs based on the semi-empirical method. *Geophysical Research Letters* 35: L02715. [doi: 10.1029/2007GL032486](https://doi.org/10.1029/2007GL032486).
- Intergovernmental Panel on Climate Change (IPCC). 1990. *IPCC First Assessment Report (FAR)*. *Climate Change: The IPCC Response Strategies*. Report prepared by Working Group III. 330 pp. http://www.ipcc.ch/publications_and_data/publications_ipcc_first_assessment_1990_wg3_shtml.
- Intergovernmental Panel on Climate Change (IPCC). 1995. *IPCC Second Assessment Report (SAR)*. *Climate Change 1995: The Science of Climate Change*. World Meteorological Organization (WMO) and United Nations Environment Programme (UNEP). 73pp. <http://www.ipcc.ch/pdf/climate-changes-1995/ipcc-2nd-assessment/2nd-assessment-en.pdf>.
- Intergovernmental Panel on Climate Change (IPCC). 2001. *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report (TAR) of the Intergovernmental Panel on Climate Change*. [JT Houghton, Y Ding, DJ Griggs, M Noguer, PJ van der Linden, X Dai, K Maskell, CA Johnson (eds.)], Cambridge University Press: Cambridge, UK, and New York, USA. 881pp. http://www.grida.no/publications/other/ipcc_tar/.
- Intergovernmental Panel on Climate Change (IPCC). 2007. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report (AR4) of the Intergovernmental Panel on Climate Change*. [S Solomon, D Qin, M Manning, M Marquis, K Averyt, MMB Tignor, HL Miller, Jr., Z Chen (eds.)], Cambridge University Press: Cambridge, UK and New York, NY, USA. 91 pp. <https://www.ipcc.ch/report/ar4/>.
- Intergovernmental Panel on Climate Change (IPCC). 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change*. [TF Stocker, D Qin, G Plattner, MMB Tignor, SK Allen, J Boschung, A Nauels, Y Xia, V Bex, PM Midgley (eds.)], Cambridge University Press: Cambridge, UK and New York, NY, USA. 1535pp. <https://www.ipcc.ch/report/ar5/>.

- Jevrejeva S, JC Moore, A Grinsted. 2008. Relative importance of mass volume changes to global sea level rise. *Journal of Geophysical Research* 113: D08105. doi: [10.1029/2007JD009208](https://doi.org/10.1029/2007JD009208).
- Jevrejeva S, JC Moore, A Grinsted. 2010. How will sea level respond to changes in natural and anthropogenic forcings by 2100? *Geophysical Research Letters* 37: L07703. doi: [10.1029/2010GL042947](https://doi.org/10.1029/2010GL042947).
- Katsman CA, GJ van Oldenborgh. 2011. Tracing the upper ocean's "missing heat". *Geophysical Research Letters* 38(14), L14610. doi: [10.1029/2011GL048417](https://doi.org/10.1029/2011GL048417).
- Konikow LF. 2011. Contribution of global groundwater depletion since 1900. *Geophysical Research Letters* 38(14): L17401. doi: [10.1029/2011GL048604](https://doi.org/10.1029/2011GL048604).
- Kopp RE, RM Horton, CM Little, JX Mitrovica, M Oppenheimer, DJ Rasmussen, BH Strauss, C Tebaldi. 2014. Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites. *Earth's Future* 2(8): 383-406. doi: [10.1002/2014EF000239](https://doi.org/10.1002/2014EF000239).
- Kopp R, F Simons, J Mitrovica, A Maloof, M Oppenheimer. 2009. Probabilistic assessment of sea level during the last interglacial stage. *Nature* 462: 863-867. doi: [10.1038/nature08686](https://doi.org/10.1038/nature08686).
- Levermann A, P Clark, B Marzeion, G Milne, D Pollard, V Radic, A Robinson. 2013. The multimillennial sea-level commitment of global warming. *Proceedings of the National Academy of Sciences* 110(34): 13745-13750. doi: [10.1073/pnas.1219414110](https://doi.org/10.1073/pnas.1219414110).
- Little CM, NM Urban, M Oppenheimer. 2013. Probabilistic framework for assessing the ice sheet contribution to sea level change. *Proceedings of the National Academy of Sciences* 110(9): 3264-3269. doi: [10.1073/pnas.1214457110](https://doi.org/10.1073/pnas.1214457110).
- Marzeion B, AH Jarosch, M Hofer. 2012. Past and future sea-level changes from the surface mass balance of glaciers. *The Cryosphere* 6: 1295-1322. doi: [10.5194/tc-6-1295-2012](https://doi.org/10.5194/tc-6-1295-2012).
- Meehl GA, WM Washington, JM Arblaster, A Hu, H Teng, C Tebaldi, BN Sanderson, J Lamarque, A Conley, WG Strand, JB White III. 2012. Climate System Response to External Forcings and Climate Change Projections in CCSM4. *Journal of Climate* 25: 3661-3683. doi: [10.1175/JCLI-D-11-00240.1](https://doi.org/10.1175/JCLI-D-11-00240.1).
- Melillo JM, TC Richmond, GW Yohe (eds). 2014. *Climate Change Impacts in the United States: The Third National Climate Assessment*. Report for the US Global Change Research Program, 841 pp. doi: [10.7930/J0Z31WJ2](https://doi.org/10.7930/J0Z31WJ2).
- National Research Council (NRC). 2012. *Sea-Level Rise for the Coasts of California, Oregon, and Washington: Past, Present, and Future*. Report by the Committee on Sea Level Rise in California, Oregon, and Washington. National Academies Press, Washington, DC. 250 pp. <http://www.nap.edu/catalog/13389/sea-level-rise-for-the-coasts-of-california-oregon-and-washington>.

- Ocean Protection Council (OPC). 2010. *Interim Guidance: Coastal and Ocean Working Group of the California Climate Action Team (CO-CAT)*.
http://opc.ca.gov/webmaster/ftp/pdf/agenda_items/20110311/12.SLR_Resolution/SLR-Guidance-Document.pdf.
- Ocean Protection Council (OPC). 2018. *State of California Sea-Level Rise Guidance: 2018 Update*.
http://www.opc.ca.gov/webmaster/ftp/pdf/agenda_items/20180314/Item3_Exhibit-A_OPC_SLR_Guidance-rd3.pdf
- Ocean Protection Council (OPC). 2013. *State of California Sea-Level Rise Guidance Document*.http://www.opc.ca.gov/webmaster/ftp/pdf/docs/2013_SLR_Guidance_Update_FINAL1.pdf.
- Overpeck JT, BL Otto-Bliesner, GH Miller, DR Muhls, RB Alley, JT Kiehl. 2006. Paleoclimatic evidence for future ice-sheet instability and rapid sea-level rise. *Science* 311(5768): 1747-1750. doi: [10.1126/science.1115159](https://doi.org/10.1126/science.1115159).
- Parris A, P Bromirski, V Burkett, D Cayan, M Culver, J Hall, R Horton, K Knuuti, R Moss, J Obeysekera, A Sallenger, J Weiss. 2012. *Global Sea Level Rise Scenarios for the US National Climate Assessment*. NOAA Tech Memo OAR CPO-1. 37 pp.
http://scenarios.globalchange.gov/sites/default/files/NOAA_SLR_r3_0.pdf.
- Perrette M, F Landerer, R Riva, K Frieler, M Meinshausen. 2013. A scaling approach to project regional sea level rise and its uncertainties. *Earth System Dynamics* 4(1): 11-29.
doi:[10.5194/esd-4-11-2013](https://doi.org/10.5194/esd-4-11-2013).
- Pfeffer WT, JT Harper, S O'Neel. 2008. Kinematic constraints on glacier contributions to 21st century sea-level rise. *Science* 321(5894): 1340 -1343. doi:[10.1126/science.1159099](https://doi.org/10.1126/science.1159099).
- Price SF, AJ Payne, IM Howat, BE Smith. 2011. Committed sea-level rise for the next century from Greenland ice sheet dynamics during the past decade. *Proceedings of the National Academy of Sciences* 108(22): 8978-8983. doi:[10.1073/pnas.1017313108](https://doi.org/10.1073/pnas.1017313108).
- Rahmstorf S. 2007. A semi-empirical approach to projecting future sea-level rise. *Science* 315(5810): 368-370. doi:[10.1126/science.1135456](https://doi.org/10.1126/science.1135456).
- Rahmstorf S, G Foster, A Cazenave. 2012. Comparing climate projections to observations up to 2011. *Environmental Research Letters* 7: 044035. doi:[10.1088/1748-9326/7/4/044035](https://doi.org/10.1088/1748-9326/7/4/044035).
- Rohling E, K Grant, C Hemleben, M Siddall, B Hoogakker, M Bolshaw, M Kucera. 2008. High rates of sea-level rise during the last interglacial period. *Nature Geoscience* 1: 38-42.
doi:[10.1038/ngeo.2007.28](https://doi.org/10.1038/ngeo.2007.28).
- Schaeffer M, W Hare, S Rahmstorf, M Vermeer. 2012. Long-term sea-level rise implied by 1.5°C and 2°C warming levels. *Nature Climate Change* 2: 867-870. doi:[10.1038/nclimate1584](https://doi.org/10.1038/nclimate1584).
- Shepherd A, ER Ivins, A Geruo, VR Barletta, MJ Bentley, S Bettadpur, KH Briggs, DH Bromwich, R Forsberg, N Galin, M Horwath, S Jacobs, I Joughin, MA King, JTM Lenaerts, J Li, SRM Ligtenberg, A Luckman, SB Luthcke, M McMillan, R Meister, G Milne, J Mouginot, A

- Muir, JP Nicolas, J Paden, AJ Payne, H Pritchard, E Rignot, H Rott, LS Sorensen, TA Scambos, B Scheuchl, EJO Schrama, B Smith, AV Sundal, JH van Angelen, WJ van de Berg, MR van den Broeke, DG Vaughan, I Velicogno, J Wahr, PL Whitehouse, DJ Wingham, D Yi, D Young, HJ Zwally. 2012. A reconciled estimate of ice-sheet mass balance. *Science* 338(6111): 1183-1189. [doi: 10.1126/science.1228102](https://doi.org/10.1126/science.1228102).
- Slangen ABA, CA Katsman, RSW van de Wal, LLA Vermeersen, REM Riva. 2012. Towards regional projections of twenty-first century sea-level change based on IPCC SRES scenarios. *Climate Dynamics* 38: 1191-1209. [doi: 10.1007/s00382-011-1057-6](https://doi.org/10.1007/s00382-011-1057-6).
- Smith WH, DT Sandwell. 1997. Global sea floor topography from satellite altimetry and ship depth soundings. *Science* 277(5334): 1956-1962. [doi: 10.1126/science.277.5334.1956](https://doi.org/10.1126/science.277.5334.1956).
- Sweet, W.V., R.E. Kopp, C.P. Weaver, J. Obeysekera, R.M. Horton, E.R. Thieler and CZ. Global and Regional Sea Level Rise Scenarios for the United States. 2017.
- Titus JG, V Narayanan. 1996. The risk of sea level rise. *Climatic Change* 33(2):151-212. [doi: 10.1007/BF00140246](https://doi.org/10.1007/BF00140246).
- Van den Broeke MR, J Bamber, J Lenaerts, E Rignot, 2011. Ice sheets and sea level: thinking outside the box. *Surveys in Geophysics* 32(4-5): 495-505. [doi: 10.1007/s10712-011-9137-z](https://doi.org/10.1007/s10712-011-9137-z).
- Vellinga P, C Katsman, A Sterl, J Beersma, W Hazeleger, J Church, R Kopp, D Kroon, M Oppenheimer, H Plag, S Rahmstorf, J Lowe, J Ridley, H von Storch, D Vaughan, R van de Wal, R Weisse, J Kwadijk, R Lammersen, N Marinova. 2009. *Exploring high-end climate change scenarios for flood protection of the Netherlands. International Scientific Assessment*, Prepared for the Delta Committee. Scientific Report WR-2009-05. KNMI, Alterra, The Netherlands. 150pp. <http://edepot.wur.nl/191831>.
- Vermeer M, S Rahmstorf. 2009. Global sea level linked to global temperature. *Proceedings of the National Academy of Science* 108: 21527-21532. [doi:10.1073/pnas.0907765106](https://doi.org/10.1073/pnas.0907765106).
- Wada Y, LPH van Beek, CM van Kempen, JWTM Reckman, S Vasak, MFP Bierkens. 2010. Global depletion of groundwater resources. *Geophysical Research Letters* 37: L20402. [doi:10.1029/2010GL044571](https://doi.org/10.1029/2010GL044571).
- Winkelman S, J Mueller, E Jue. 2012. *Climate Adaptation & Transportation: Identifying Information and Assistance Needs*. Summary of an Expert Workshop held November 2011, Center for Clean Air Policy & Environmental and Energy Study Institute. http://ccap.org/assets/Climate-Adaptation-and-Transportation_CCAP-EESI-May-2012.pdf.
- Yin J, SM Griffies, RJ Stouffer. 2010. Spatial variability of sea-level rise in the twenty-first century projections. *Journal of Climate* 23(17): 4585-4607. [doi: 10.1175/2010JCLI3533.1](https://doi.org/10.1175/2010JCLI3533.1).
- Zecca A, L Chiari. 2012. Lower bounds to future sea-level rise. *Global and Planetary Change* 98-99: 1-5. [doi:10.1016/j.gloplacha.2012.08.002](https://doi.org/10.1016/j.gloplacha.2012.08.002).