

CALIFORNIA COASTAL COMMISSION

SOUTH CENTRAL COAST AREA
89 SOUTH CALIFORNIA ST., SUITE 200
VENTURA, CA 93001
(805) 585-1800

Filed: 3/14/08
180th Day: 9/10/08
Staff: Shana Gray
Staff Report: 5/28/08
Hearing Date: 6/11/08



W17a

STAFF REPORT: REGULAR CALENDAR

APPLICATION NO.: 4-06-153

APPLICANT: Los Angeles County Department of Public Works

PROJECT DESCRIPTION: Construction of a 71 linear foot soldier pile retaining wall, a maximum of 4 feet 6 inches above road level, consisting of a 2-ft. 3-inch handrail on top of a 2-ft. 3-inch concrete barrier rail with colored concrete panels. The project includes reconstruction of 795 sq. ft. of the Grandview Drive roadway; removal of existing guard rail, chain link fence and one mature pine tree; temporary k-rail during construction; revegetation of the 0.05-acre temporary construction easement with native plant species after construction of the retaining wall has been completed, and 187 cu. yds. (129 cu. yds. cut, 58 cu. yds fill) of grading.

The project further includes after-the-fact authorization of: (1) a 90 linear ft soldier pile retaining wall comprised of a 2-ft. 3-inch concrete barrier rail with chain link fence on top to be retrofitted with 2-ft. 3-inch handrail to match proposed wall, (2) reconstruction of approximately 1,210 sq. ft. of Grandview Drive roadway, (3) revegetation of the 0.07 acre impact area, and (4) and 89 cu. yds. (56 cu. yds cut, and 33 cu. yds. fill) of grading.

PROJECT LOCATION: Grandview Drive, 60 feet North of Falls Drive, Fernwood Small Lot Subdivision, in the Santa Monica Mountains; Los Angeles County (APNs 4447-002-008; 4447-002-010).

MOTION & RESOLUTION: Page 4

SUMMARY OF STAFF RECOMMENDATION:

Staff recommends **APPROVAL** of the proposed development with seven special conditions regarding: (1) a restoration plan; (2) oak tree mitigation and monitoring; (3) assumption of risk; (4) material and design specification; (5) removal of excess excavated material; (6) best management practices; and (7) condition compliance.

Construction of the new wall and reconstruction of the roadway are proposed to remediate an outboard slope failure approximately 66 feet long which resulted from the January 2005 storm events, and caused visible damage to the roadway. The applicant has stated that the 1997 construction of the 90 linear-ft. wall and reconstruction of the roadway were necessary to remediate slope failure and road damage that occurred during the 1995 storm season. The new wall would be a continuation of the south end of the existing unpermitted wall, and upon project completion, would appear as one integrated design. The purpose of the proposed project is to maintain the public's ability to use Grandview Drive for vehicular access and provide for emergency services/access to the residences and communities along this roadway.

The project site is located within oak woodland habitat, a habitat that is considered to be environmentally sensitive habitat area (ESHA) in the Santa Monica Mountains. The project footprint is mostly disturbed due to its location adjacent to the road shoulder and because the outboard slope is located adjacent to existing residential development that was constructed prior to the Coastal Act. As a result, although the oak canopies extend over a portion of the project site, the understory of the oak trees is significantly disturbed, supporting scattered ornamental plantings. However, the proposed project will not involve, or require, the removal of any oak trees on site.

The County has submitted an engineering and alternatives analysis which finds that the proposed soldier pile retaining wall is necessary to stabilize the downslope area of Grandview Drive in order to prevent further slope failure that could undermine the public roadway. Given that slope protection is needed in order to prevent erosion of the road shoulder in a fixed position that is constrained by the location of sensitive habitat, it is not possible to redesign or relocate the proposed development in a manner that would provide for a buffer from the sensitive habitat areas on site.

Therefore design alternatives, rather than siting alternatives, were the focus of the alternatives analysis. The design alternatives that were considered include the removal and recompaction of fill to regrade the slope and the construction of a concrete retaining wall. However, both of these design options would require a larger project footprint resulting in significant vegetation removal and impacts to ESHA. In addition, the "no project", or road closure, alternative is not viable in this case because Grandview Drive is a public roadway that must be maintained for vehicular and emergency access to the Fernwood subdivision. In addition, the "no project" alternative would result in continued slope failure and erosion.

Though the proposed soldier pile wall introduces a large physical structure into the project area, this alternative would minimize potential impacts to ESHA, would revegetate the downslope area, and would be designed to minimize visual intrusion into the surrounding environment to the maximum extent feasible. Staff concurs that there are no less environmentally damaging alternatives to the proposed project that would avoid or further reduce impacts to sensitive coastal resources.

However, in order to ensure that adjacent ESHA is protected, special conditions been included to require the applicant to: implement a oak woodland habitat restoration plan that includes revegetation with native vegetation for all areas that are temporarily disturbed by grading and construction activities; mitigate for the significant encroachment into three oak tree canopy driplines; remove all excess fill material from the project site; and implement best management practices to minimize erosion and protect coastal water quality.

The standard of review for the proposed permit application is the Chapter Three policies of the Coastal Act. In addition, the policies of the certified Malibu – Santa Monica Mountains Land Use Plan (LUP) serve as guidance. The proposed project, as conditioned, employs a method that is consistent with the applicable resource protection provisions of the Coastal Act.

TABLE OF CONTENTS

I. STAFF RECOMMENDATION.....	4
II. STANDARD CONDITIONS.....	4
III. SPECIAL CONDITIONS	5
1. <i>Restoration Plan</i>	5
2. <i>Oak Tree Mitigation and Monitoring</i>	7
3. <i>Assumption of Risk</i>	7
4. <i>Material/Design Specifications</i>	8
5. <i>Removal of Excess Excavated Material</i>	8
6. <i>Best Management Practices</i>	8
7. <i>Condition Compliance</i>	8
IV. FINDINGS AND DECLARATIONS	9
A. PROJECT DESCRIPTION AND BACKGROUND	9
B. ENVIRONMENTALLY SENSITIVE HABITAT AREAS AND WATER QUALITY	10
1. <i>ESHA Determination</i>	11
2. <i>Application of the Section 30240 ESHA Protection Policy</i>	15
3. <i>Water Quality</i>	18
C. HAZARDS AND GEOLOGIC STABILITY	19
D. VISUAL RESOURCES	21
E. UNPERMITTED DEVELOPMENT	22
F. LOCAL COASTAL PROGRAM.....	23
G. CEQA.....	23

EXHIBITS

- Exhibit 1. Vicinity Map**
Exhibit 2. Site Plan

Exhibit 3. Wall Plan

Exhibit 4. Railing Plan

Exhibit 5. Dr. John Dixon Memorandum Regarding Environmentally Sensitive
Habitat Areas In The Santa Monica Mountains

LOCAL APPROVALS RECEIVED: N/A

SUBSTANTIVE FILE DOCUMENTS: 2006 Biological Reconnaissance Survey Results for the Grandview Drive Repair Project (URS Corporation, September 2006); Engineering Memo for Proposed Retaining Wall on Grandview Drive (Los Angeles County Department of Public Works, no date, received in Commission Office December 14, 2007); Caltrans Memo to Bridge Designers, Section 13-1; Caltrans Highway Design Manual, page 200-47;

I. STAFF RECOMMENDATION

MOTION: *I move that the Commission approve Coastal Development Permit No. 4-06-153 pursuant to the staff recommendation.*

STAFF RECOMMENDATION OF APPROVAL:

Staff recommends a **YES** vote. Passage of this motion will result in approval of the permit as conditioned and adoption of the following resolution and findings. The motion passes only by affirmative vote of a majority of the Commissioners present.

RESOLUTION TO APPROVE THE PERMIT:

The Commission hereby approves a coastal development permit for the proposed development and adopts the findings set forth below on grounds that the development as conditioned will be in conformity with the policies of Chapter 3 of the Coastal Act and will not prejudice the ability of the local government having jurisdiction over the area to prepare a Local Coastal Program conforming to the provisions of Chapter 3. Approval of the permit complies with the California Environmental Quality Act because either 1) feasible mitigation measures and/or alternatives have been incorporated to substantially lessen any significant adverse effects of the development on the environment, or 2) there are no further feasible mitigation measures or alternatives that would substantially lessen any significant adverse impacts of the development on the environment.

II. STANDARD CONDITIONS

1. Notice of Receipt and Acknowledgment. The permit is not valid and development shall not commence until a copy of the permit, signed by the permittee or authorized agent,

acknowledging receipt of the permit and acceptance of the terms and conditions, is returned to the Commission office.

2. **Expiration.** If development has not commenced, the permit will expire two years from the date on which the Commission voted on the application. Development shall be pursued in a diligent manner and completed in a reasonable period of time. Application for extension of the permit must be made prior to the expiration date.
3. **Interpretation.** Any questions of intent or interpretation of any condition will be resolved by the Executive Director or the Commission.
4. **Assignment.** The permit may be assigned to any qualified person, provided assignee files with the Commission an affidavit accepting all terms and conditions of the permit.
5. **Terms and Conditions Run with the Land.** These terms and conditions shall be perpetual, and it is the intention of the Commission and the permittee to bind all future owners and possessors of the subject property to the terms and conditions.

III. SPECIAL CONDITIONS

1. Restoration Plan

Prior to issuance of this Coastal Development Permit, the applicant shall submit, for the review and approval of the Executive Director, a detailed Restoration and Monitoring Plan, prepared by a biologist or environmental resource specialist with qualifications acceptable to the Executive Director, for all disturbed areas within the temporary construction easement and any other areas of the project site temporarily disturbed by grading and construction activities. Within 90 days of the completion of construction, the applicant shall commence implementation of the approved Restoration and Monitoring Plan. The Executive Director may grant additional time for good cause. The plans shall identify the species, extent, and location of all plant materials to be removed or planted and shall incorporate the following criteria:

a. Technical Specifications

The Restoration and Monitoring Plan shall provide for the restoration of the oak woodland habitat understory in the project area with native plant species that are appropriate for oak woodlands, including coverage of areas downslope of the 71 feet of the proposed wall and where vegetation has been temporarily disturbed or removed due to construction activities. These areas shall be replanted with native plant species that are appropriate for oak woodland habitat in the same general location. The restoration area shall be delineated on a site plan. All invasive and non-native plant species shall be removed from the restoration area.

The plan shall include detailed documentation of conditions on site prior to the approved construction activity (including photographs taken from pre-designated sites annotated

to a copy of the site plans) and specify restoration goals and specific performance standards to judge the success of the restoration effort.

The plan shall also provide information on removal methods for exotic species, salvage of existing vegetation, revegetation methods and vegetation maintenance. The plan shall further include details regarding the types, sizes, and location of plants to be placed within the restoration area. Only native plant species appropriate for a southern oak woodland and which are endemic to the Santa Monica Mountains shall be used, as listed by the California Native Plant Society - Santa Monica Mountains Chapter in their document entitled Recommended List of Plants for Landscaping in the Santa Monica Mountains, dated February 5, 1996. All native plant species shall be of local genetic stock. No plant species listed as problematic and/or invasive by the California Native Plant Society, the California Invasive Plant Council, or by the State of California shall be employed or allowed to naturalize or persist on the site. No plant species listed as a 'noxious weed' by the State of California or the U.S. Federal Government shall be utilized or maintained within the property. Site restoration shall be deemed successful if the revegetation of native plant species on site is adequate to provide 90% coverage by the end of the five (5) year monitoring period and is able to survive without additional outside inputs, such as supplemental irrigation. The plan shall also include a detailed description of the process, materials, and methods to be used to meet the approved goals and performance standards and specify the preferable time of year to carry out restoration activities and describe the interim supplemental watering requirements that will be necessary.

b. Monitoring Program

A monitoring program shall be implemented to monitor the project for compliance with the specified guidelines and performance standards. The applicant shall submit, upon completion of the initial planting, a written report prepared by a qualified resource specialist, for the review and approval of the Executive Director, documenting the completion of the initial planting/revegetation work. This report shall also include photographs taken from pre-designated sites (annotated to a copy of the site plans) documenting the completion of the initial planting/revegetation work.

Five years from the date of issuance of this coastal development permit, the applicant shall submit for the review and approval of the Executive Director, a Restoration Monitoring Report, prepared by a qualified biologist or Resource Specialist, which certifies whether the on-site restoration is in conformance with the restoration plan approved pursuant to this Special Condition. The monitoring report shall include photographic documentation of plant species and plant coverage.

If the monitoring report indicates the restoration is not in conformance with or has failed to meet the performance standards specified in the restoration plan approved pursuant to this permit, the applicant, or successors in interest, shall submit a revised or supplemental restoration plan for the review and approval of the Executive Director and shall implement the approved version of the plan. The revised restoration plan must be

prepared by a qualified biologist or Resource Specialist and shall specify measures to remediate those portions of the original plan that have failed or are not in conformance with the original approved plan.

2. Oak Tree Mitigation and Monitoring

A. Prior to issuance of the Coastal Development Permit, the applicant shall submit, for the review and approval of the Executive Director, an off-site oak tree replacement planting program, which specifies replacement tree locations, tree or seedling size planting specifications, and a ten-year monitoring program with specific performance standards to ensure that the replacement planting program is successful. The applicant may coordinate with the Mountains Restoration Trust, or other entity acceptable to the Executive Director, to prepare and implement the off-site oak tree replacement planting program. At least 30 replacement seedlings, less than one year old, grown from acorns collected in the selected planting area, shall be planted as mitigation for impacts to three oak trees as a result of the proposed project. All replacement seedlings must be planted in a suitable location that is restricted in perpetuity from development or is public parkland. An appropriate off-site planting area may be identified in collaboration with the Mountains Restoration Trust (MRT), or other entity acceptable to the Executive Director, and shall be located in the Topanga Canyon watershed within the coastal zone of the Santa Monica Mountains.

B. An annual monitoring report on the oak tree replacement area shall be submitted for the review and approval of the Executive Director for each of the 10 years. If monitoring indicates the oak trees are not in conformance with or has failed to meet the performance standards specified in the monitoring program approved pursuant to this permit, the applicant, or successors in interest, shall submit a revised or supplemental planting plan for the review and approval of the Executive Director. The revised planting plan shall specify measures to remediate those portions of the original plan that have failed or are not in conformance with the original approved plan.

C. The applicant shall commence implementation of the approved off-site oak tree replacement planting program concurrently with the commencement of construction on the project site. The Executive Director may extend this time for good reason.

3. Assumption of Risk

By acceptance of this permit, the applicant acknowledges and agrees (i) that the site may be subject to hazards from erosion, landslide, and slope failure; (ii) to assume the risks to the applicant and the property that is the subject of this permit of injury and damage from such hazards in connection with this permitted development; (iii) to unconditionally waive any claim of damage or liability against the Commission, its officers, agents, and employees for injury or damage from such hazards; and (iv) to indemnify and hold harmless the Commission, its officers, agents, and employees with respect to the Commission's approval of the project against any and all liability, claims, demands, damages, costs (including costs and fees incurred in defense of such claims),

expenses, and amounts paid in settlement arising from any injury or damage due to such hazards.

4. Material/Design Specifications

Prior to issuance of this Coastal Development Permit, the applicant shall submit detailed plans, for the review and approval of the Executive Director, which show that all exposed surfaces of the approved soldier pile retaining wall and handrail, shall be designed to include, blend, or mimic, the native materials and appearance (including color and texture) of the natural environment (such as the appearance of rock facing). The existing 90 linear feet of chain link fencing on the existing wall shall be replaced with the proposed metal handrail, matching with, and blending to, the handrail on the adjacent 70 linear ft new soldier pile wall.

5. Removal of Excess Excavated Material

Prior to issuance of this Coastal Development Permit, the applicant shall provide evidence to the Executive Director of the location of the disposal site for all excess excavated material from the site. If the disposal site is located in the Coastal Zone, the disposal site must have a valid coastal development permit for the disposal of fill material. If the disposal site is not authorized to receive fill material and/or does not have a coastal permit, such a permit will be required prior to the disposal of material.

6. Best Management Practices.

By acceptance of this permit, the applicant agrees to implement best management practices to minimize erosion and protect water quality adjacent to the project site. These measures shall include minimizing grading during the rainy season (November 1 – March 31). Should construction activities occur during the rainy season, the applicant shall implement sediment control measures such as sand bag barriers, silt fencing, geofabric covers or other appropriate cover, or geotextiles and/or mats on all cut or fill slopes as soon as possible. These erosion measures shall be required on the project site prior to or concurrent with the initial grading operations and maintained throughout the development process to minimize erosion and sediment from runoff waters during construction. Furthermore, should grading or site preparation cease for a period of more than 30 days, temporary erosion control measures shall be implemented and should include, but not be limited to: stabilization of all stockpiled fill, disturbed soils and cut and fill slopes with geotextiles and/or mats, sand bag barriers, and silt fencing. These temporary erosion control measures shall be monitored and maintained until grading or construction operations resume.

7. Condition Compliance

Within 120 days of Commission action on this coastal development permit application, or within such additional time as the Executive Director may grant for good cause, the applicant shall satisfy all requirements specified in the conditions hereto that the applicant is required to satisfy prior to issuance of this permit. Failure to comply with

this requirement may result in the institution of enforcement action under the provisions of Chapter 9 of the Coastal Act.

IV. FINDINGS AND DECLARATIONS

The Commission hereby finds and declares:

A. PROJECT DESCRIPTION AND BACKGROUND

The subject site is located on Grandview Drive, approximately 60 feet north of its intersection with Falls Drive, within the Fernwood small lot subdivision community, in the Santa Monica Mountains; Los Angeles County (Exhibit 1). The project crosses two privately-owned parcels. The property owners have granted permission through a temporary construction easement for L.A. County Public Works to access the subject property and complete the proposed project.

The project site drains to a blueline stream that runs along Falls Drive, under Topanga Canyon Boulevard and drains into Topanga Canyon Creek approximately 500 feet from the subject site.

The proposed project consists of construction of a 71 linear foot soldier pile retaining wall, a maximum of 4 feet 6 inches above road level, consisting of a 2-ft. 3-inch handrail on top of a 2-ft. 3-inch concrete barrier rail with colored concrete panels. The project also includes reconstruction of 795 sq. ft. of the Grandview Drive roadway; removal of existing guard rail, chain link fence and one mature pine tree; temporary k-rail during construction; revegetation of the 0.05-acre temporary construction easement with native plant species after construction of the retaining wall has been completed, and 187 cu. yds. (129 cu. yds. cut, 58 cu. yds. fill) of grading. Construction of the wall is proposed to remediate an outboard slope failure approximately 66 feet long which resulted from the January 2005 storm events, and caused visible damage to the roadway. (Exhibits 2 – 4)

The subject application further requests after-the-fact authorization for a soldier pile wall previously constructed along Grandview Drive without the benefit of a coastal development permit. The wall was constructed in 1997 by Los Angeles County Public Works Department to restore the road after the 1995 storms caused slope failure below the road and damaged the roadway. The proposed 71 ft. wall would be a continuation of the south end of the existing unpermitted wall, and upon project completion, would appear as one integrated design. This application includes a request for after-the-fact authorization for: (1) a 90 linear ft soldier pile retaining wall comprised of a 2-ft. 3-inch concrete barrier rail with chain link fence on top to be retrofitted with 2-ft. 3-inch handrail to match proposed wall, (2) reconstruction of approximately 1,210 sq. ft. of Grandview Drive roadway, (3) revegetation of the 0.07 acre impact area, and (4) and 89 cu. yds. (56 cu. yds. cut, and 33 cu. yds. fill) of grading.

Coastal Permit Required for Repair and Maintenance within ESHA

The proposed work is designed to repair a damaged public roadway. The project entails repair and maintenance work. The Commission has expressly recognized, since 1978, certain types of repair and maintenance work related to roads as exempt from permit requirements pursuant to Section 13252 of the Commission's regulations and Section 30610(d) of the Public Resource Code. See California Public Resources Code ("PRC") Section 30610(d) and the "Repair, Maintenance and Utility Hook-Up Exclusions From Permit Requirements" (adopted by the Commission on Sept. 5, 1978) (hereafter, "R&M Exclusions") Appendix I, § 3 (referring to "installation of slope protection devices, minor drainage facilities"). However, the exemptions provided by the above referenced sections and the R&M Exclusions are limited. Accordingly, California Code of Regulations, Title 14 ("14 CCR"), Section 13252 (a) lists extraordinary methods of repair and maintenance that do still require a permit. Among those methods is any repair or maintenance "located in an environmentally sensitive habitat area." 14 CCR § 13252(a)(3). Since this project would occur within such an area, the method by which this project is conducted is not exempt, and a permit is required. In addition, further review of the R&M Exclusions Guidelines confirms that this proposed repair and maintenance is not exempt from permit requirements based on that document because the proposed development is located outside the "roadway prism" or the roadway property or easement and includes construction of a new soldier pile retaining wall.

Similarly, 14 CCR Section 13252(a) states that "activities specifically described in the [R&M Exclusions guidance document that] that will have a risk of substantial adverse impact on . . . environmentally sensitive habitat area" are not exempt based on that document and may require a coastal development permit, pursuant to the normal application of section 13252. Thus, in this case, although the project is a repair and maintenance project, since the work is to be performed within an ESHA, Section 13252(a)'s limits on the repair and maintenance exemption do apply, and this project does require a permit to ensure that the method employed is as consistent as possible with the Chapter 3 policies of the Coastal Act. Moreover, this project involves excavation, and the R&M Exclusions guidance document expressly states that a permit is required "for excavation . . . outside of the roadway prism" Id. at § II.A., page 2. Therefore, a coastal development permit is required for this project.

B. ENVIRONMENTALLY SENSITIVE HABITAT AREAS AND WATER QUALITY

Section 30231 of the Coastal Act states:

The biological productivity and the quality of coastal waters, streams, wetlands, estuaries, and lakes appropriate to maintain optimum populations of marine organisms and for the protection of human health shall be maintained and, where feasible, restored through, among other means, minimizing adverse effects of waste water discharges and entrainment, controlling runoff, preventing depletion of ground water supplies and substantial interference with surface water flow, encouraging waste water

reclamation, maintaining natural vegetation buffer areas that protect riparian habitats, and minimizing alteration of natural streams.

Section 30240(b) of the Coastal Act states:

Development in areas adjacent to environmentally sensitive habitat areas and parks and recreation areas shall be sited and designed to prevent impacts which would significantly degrade those areas, and shall be compatible with the continuance of those habitat and recreation areas.

Section 30107.5 of the Coastal Act, defines an environmentally sensitive area as:

"Environmentally sensitive area" means any area in which plant or animal life or their habitats are either rare or especially valuable because of their special nature or role in an ecosystem and which could be easily disturbed or degraded by human activities and developments.

Section 30231 of the Coastal Act requires that the biological productivity and the quality of coastal waters and streams be maintained and, where feasible, restored through among other means, minimizing adverse effects of waste water discharge and entrainment, controlling runoff, preventing depletion of ground water supplies and substantial interference with surface water flows, maintaining natural buffer areas that protect riparian habitats, and minimizing alteration of natural streams. In addition, Section 30240 of the Coastal Act states that environmentally sensitive habitat areas must be protected against disruption of habitat values.

The project includes reconstruction of an existing developed roadway and the construction of a new soldier pile retaining wall; after-the-fact authorization for the construction of a soldier pile retaining wall and reconstruction of the roadway along Grandview Drive; and revegetation of the temporary construction easement with native plant species after construction of the retaining wall has been completed. One pine tree is proposed to be removed.

1. ESHA Determination

According to the applicant's submitted biological reconnaissance survey (URS Corporation, 2006), the project site is adjacent to and surrounded by residences within a mosaic of sparse native oak woodland. Staff has consulted aerial photographs of the surrounding area and has confirmed that the project site is adjacent to larger, contiguous areas of sensitive habitat.

The project area is located along Grandview Drive, immediately upslope and adjacent to existing residential development that was constructed prior to the implementation of the Coastal Act. The project footprint is mostly disturbed due to its location adjacent to the road shoulder and due to its proximity to the adjacent residential development. However, although the understory is disturbed by existing development and associated ornamental plantings, the project site is also located underneath native coast live oak woodland canopy. Oak tree canopies extend over a portion of the project site, emanating from a limited number of trees situated adjacent to the project site, including

trees that are physically located on the opposite side of Grandview Drive. No oak trees would be removed; however, the project would encroach within the driplines of three oak trees that are located immediately adjacent to the project area.

The Santa Monica Mountains comprise the largest, most pristine, and ecologically complex example of a Mediterranean ecosystem in coastal southern California. California's coastal sage scrub, chaparral, oak woodlands, and associated riparian areas have analogues in just a few areas of the world with similar climate. Mediterranean ecosystems with their wet winters and warm dry summers are only found in five localities (the Mediterranean coast, California, Chile, South Africa, and south and southwest Australia). Throughout the world, this ecosystem with its specially adapted vegetation and wildlife has suffered severe loss and degradation from human development. Worldwide, only 18 percent of the Mediterranean community type remains undisturbed¹. However, within the Santa Monica Mountains, this ecosystem is remarkably intact despite the fact that it is closely surrounded by some 17 million people. For example, the 150,000 acres of the Santa Monica Mountains National Recreation Area, which encompasses most of the Santa Monica Mountains, was estimated to be 90 percent free of development in 2000². Therefore, this relatively pristine area is both large and mostly unfragmented, which fulfills a fundamental tenet of conservation biology³. The need for large contiguous areas of natural habitat in order to maintain critical ecological processes has been emphasized by many conservation biologists⁴.

The habitat integrity and connectivity that is still evident within the Santa Monica Mountains is extremely important to maintain, because both theory and experiments over 75 years in ecology confirm that large spatially connected habitats tend to be more stable and have less frequent extinctions than habitats without extended spatial

¹ National Park Service. 2000. Draft general management plan & environmental impact statement. Santa Monica Mountains National Recreation Area – California.

² Ibid.

³ Harris, L. D. 1988. Edge effects and conservation of biotic diversity. *Conserv. Biol.* 330-332. Soule, M. E, D. T. Bolger, A. C. Alberts, J. Wright, M. Sorice and S. Hill. 1988. Reconstructed dynamics of rapid extinctions of chaparral-requiring birds in urban habitat islands. *Conserv. Biol.* 2: 75-92. Yahner, R. H. 1988. Changes in wildlife communities near edges. *Conserv. Biol.* 2:333-339. Murphy, D. D. 1989. Conservation and confusion: Wrong species, wrong scale, wrong conclusions. *Conservation Biol.* 3:82-84.

⁴ Crooks, K. 2000. Mammalian carnivores as target species for conservation in Southern California. p. 105-112 *in*: Keeley, J. E., M. Baer-Keeley and C. J. Fotheringham (eds), 2nd Interface Between Ecology and Land Development in California, U.S. Geological Survey Open-File Report 00-62. Sauvajot, R. M., E. C. York, T. K. Fuller, H. Sharon Kim, D. A. Kamradt and R. K. Wayne. 2000. Distribution and status of carnivores in the Santa Monica Mountains, California: Preliminary results from radio telemetry and remote camera surveys. p 113-123 *in*: Keeley, J. E., M. Baer-Keeley and C. J. Fotheringham (eds), 2nd Interface Between Ecology and Land Development in California, U.S. Geological Survey Open-File Report 00-62. Beier, P. and R. F. Noss. 1998. Do habitat corridors provide connectivity? *Conserv. Biol.* 12:1241-1252. Beier, P. 1996. Metapopulation models, tenacious tracking and cougar conservation. *In*: Metapopulations and Wildlife Conservation, ed. D. R. McCullough. Island Press, Covelo, California, 429p.

structure⁵. Beyond simply destabilizing the ecosystem, fragmentation and disturbance can even cause unexpected and irreversible changes to new and completely different kinds of ecosystems (habitat conversion)⁶.

As a result of the pristine nature of large areas of the Santa Monica Mountains and the existence of large, unfragmented and interconnected blocks of habitat, this ecosystem continues to support an extremely diverse flora and fauna. The observed diversity is probably a function of the diversity of physical habitats. The Santa Monica Mountains have the greatest geological diversity of all major mountain ranges within the transverse range province. According to the National Park Service, the Santa Monica Mountains contain 40 separate watersheds and over 170 major streams with 49 coastal outlets⁷. These streams are somewhat unique along the California coast because of their topographic setting. As a “transverse” range, the Santa Monica Mountains are oriented in an east-west direction. As a result, the south-facing riparian habitats have more variable sun exposure than the east-west riparian corridors of other sections of the coast. This creates a more diverse moisture environment and contributes to the higher biodiversity of the region. The many different physical habitats of the Santa Monica Mountains support at least 17 native vegetation types⁸ including the following habitats considered sensitive by the California Department of Fish and Game: native perennial grassland, coastal sage scrub, red-shank chaparral, valley oak woodland, walnut woodland, southern willow scrub, southern cottonwood-willow riparian forest, sycamore-alder woodland, oak riparian forest, coastal salt marsh, and freshwater marsh. Over 400 species of birds, 35 species of reptiles and amphibians, and more than 40 species of mammals have been documented in this diverse ecosystem. More than 80 sensitive species of plants and animals (listed, proposed for listing, or species of concern) are known to occur or have the potential to occur within the Santa Monica Mountains Mediterranean ecosystem.

The project is located within coast live oak woodland habitat, a habitat that is considered to be environmentally sensitive habitat area (ESHA) in the Santa Monica Mountains. As described above and in more detail in the memo “Designation of ESHA in the Santa Monica Mountains, dated March 25, 2003 by John Dixon (Exhibit 5), the Mediterranean Ecosystem in the Santa Mountains is itself both rare and valuable

⁵ Gause, G. F. 1934. The struggle for existence. Baltimore, William and Wilkins 163 p. (also reprinted by Hafner, N.Y. 1964). Gause, G. F., N. P. Smaragdova and A. A. Witt. 1936. Further studies of interaction between predators and their prey. J. Anim. Ecol. 5:1-18. Huffaker, C. B. 1958. Experimental studies on predation: dispersion factors and predator-prey oscillations. Hilgardia 27:343-383. Luckinbill, L. S. 1973. Coexistence in laboratory populations of *Paramecium aurelia* and its predator *Didinium nasutum*. Ecology 54:1320-1327. Allen, J. C., C. C. Brewster and D. H. Slone. 2001. Spatially explicit ecological models: A spatial convolution approach. Chaos, Solitons and Fractals. 12:333-347.

⁶ Scheffer, M., S. Carpenter, J. A. Foley, C. Folke and B. Walker. 2001. Catastrophic shifts in ecosystems. Nature 413:591-596.

⁷ NPS. 2000. op.cit.

⁸ From the NPS report (2000 op. cit.) that is based on the older Holland system of subjective classification. The data-driven system of Sawyer and Keeler-Wolf results in a much larger number of distinct “alliances” or vegetation types.

because of its relatively pristine character, physical complexity, and resultant biological diversity. Oak woodland and savannahs are a component of this Mediterranean Ecosystem. Coast live oak woodland occurs mostly on north slopes, shaded ravines and canyon bottoms. Besides the coast live oak, this plant community includes hollyleaf cherry, California bay laurel, coffeeberry, and poison oak. Coast live oak also occurs as a riparian corridor species within the Santa Monica Mountains.

In the Santa Monica Mountains, oak woodlands provide habitat that has many important roles in the ecosystem, including the provision of critical linkages to other community types such as riparian corridors and grassland, the provision of essential habitat for a high diversity of birds, the provision of refuge for raptors and bats, the support of rare species, and the reduction of erosion on slopes, thereby protecting the water quality of coastal streams. Typical wildlife in this habitat includes acorn woodpeckers, scrub jays, plain titmice, northern flickers, cooper's hawks, western screech owls, mule deer, gray foxes, ground squirrels, jackrabbits and several species of sensitive bats.

The natural habitats of the Santa Monica Mountains, including oak woodland communities, are highly threatened by current development pressure, fragmentation and impacts from the surrounding megalopolis. The developed portions of the Santa Monica Mountains represents the extension of this urbanization into natural areas. About 54% of the undeveloped Santa Monica Mountains are in private ownership⁹, and computer simulation studies of the development patterns over the next 25 years predict a serious increase in habitat fragmentation¹⁰. Development and associated human activities have many well-documented deleterious effects on natural communities. These environmental impacts may be both direct and indirect and include the effects of increased fire frequency, of fire clearance, of introduction of exotic species, and of night lighting.

Therefore, for the reasons detailed above and further detailed in Exhibit 5, which is incorporated herein, due to its important ecosystem functions and vulnerability to development, oak woodland habitat within the Santa Monica Mountains meet the definition of ESHA under the Coastal Act. Further, though the understory is disturbed, the Commission finds that the oak woodland at the subject site meets the definition of ESHA, pursuant to Section 30107.5 of the Coastal Act, due to: its increasing rarity, its connectivity with larger contiguous habitats, its vital role in the functioning of the Santa Monica Mountains Mediterranean ecosystem, its important ecosystem functions, and its vulnerability to development.

⁹ National Park Service. 2000. Draft: General Management Plan & Environmental Impact Statement, Santa Monica Mountains National Recreation Area, US Dept. of Interior, National Park Service, December 2000.

¹⁰ Swenson, J. J., and J. Franklin. 2000. The effects of future urban development on habitat fragmentation in the Santa Monica Mountains. *Landscape Ecol.* 15:713-730.

2. Application of the Section 30240 ESHA Protection Policy

Coastal Act Section 30240 requires that ESHA be protected against any significant disruption of habitat values, and that only resource dependent uses be allowed within ESHA. Further, Section 30240 requires that development in areas adjacent to environmentally sensitive habitat areas be sited and designed to prevent impacts which would significantly degrade such areas and shall be compatible with the continuance of such habitat areas. In addition, Section 30231 of the Coastal Act specifically provides that the quality of coastal waters and streams shall be maintained and restored whenever feasible.

To assist in the determination of whether a project is consistent with Sections 30231 and 30240 of the Coastal Act, the Commission has, in past coastal development permit actions for new development in the Santa Monica Mountains, looked to the certified Malibu/Santa Monica Mountains Land Use Plan (LUP) for guidance. The 1986 LUP has been found to be consistent with the Coastal Act and provides specific standards for development within the Santa Monica Mountains. In its findings regarding the certification of the Malibu/Santa Monica Mountains LUP, the Commission emphasized the importance placed by the Coastal Act on protection of sensitive environmental resources finding that:

Environmentally sensitive habitat areas (ESHAs) shall be protected against significant disruption of habitat values, and only uses dependent on such resources shall be allowed within such areas. Residential use shall not be considered a resource dependent use.

Specifically, Policy 76 of the LUP, in concert with the Coastal Act, limits stream alterations. In addition, Policy 82 of the LUP, in concert with the Coastal Act, provides that grading shall be minimized to ensure that the potential negative effects of runoff and erosion on watershed and streams is minimized. Further, Policies 84 and 94, in concert with the Coastal Act, provide that disturbed areas shall be revegetated with native plant species within environmentally sensitive habitat areas and significant watersheds.

In this case, the proposed road reconstruction and soldier pile retaining wall is necessary to stabilize the existing roadway that was damaged by soil erosion associated with a large storm event. Given that the project addresses a slope failure in a fixed location, the siting of such development to avoid impacts to ESHA is necessarily constrained. In this case, the failing slope itself is located partially underneath oak woodland ESHA. As a result, it is not possible to relocate the proposed development in a manner that would avoid or provide a buffer from the sensitive habitat areas. Therefore, it is essential to consider design options that would reduce impacts to ESHA, consistent with Coastal Act Section 30240.

The proposed project is designed to repair the existing public road that was previously damaged due to storm activity. The project entails necessary repair and maintenance work. As discussed in detail above, although the proposed repair of the existing public

roadway and its supporting slopes is generally consistent with the repair and maintenance activities that are allowed under Coastal Act and the R&M Guidelines for public projects, in this case, the project is located within ESHA and therefore, a coastal development permit is required.

The County Geotechnical and Materials Engineering Division has submitted an engineering and alternatives analysis for the proposed project, which indicates that the excessive runoff during the January 2005 storm season caused the road and hillside to fail along this section of Grandview Drive. The report noted that the large cracks in the roadway are due to “soil slumping caused by intense rainfall and ensuing saturation of the slope” and that “without construction of the soldier pile retaining wall the un-braced slope is susceptible to failure in the next occurrence of intense rainfall. Failure of the slope will render the roadway impassable and pose a public safety hazard.”

The engineering and alternatives analysis found that the proposed soldier pile retaining wall is necessary to stabilize the downslope area of Grandview Drive in order to prevent further slope failure that could undermine the public roadway. The analysis indicates that design alternatives that were considered include the removal and recompaction of fill to regrade the slope and the construction of a concrete retaining wall. However, both of these design options would require a larger project footprint resulting in significant vegetation removal and impacts to ESHA. In addition, the “no project”, or road closure, alternative is not viable in this case because Grandview Drive is a public roadway that must be maintained for vehicular and emergency access to the Fernwood subdivision. In addition, the “no project” alternative would result in continued slope failure and erosion.

Though the proposed soldier pile wall introduces a large physical structure into the project area, this alternative would minimize potential impacts to ESHA, would revegetate the downslope area, and would be designed to minimize visual intrusion into the surrounding environment. Staff concurs that there are no less environmentally damaging alternatives to the proposed project that would avoid or further reduce impacts to sensitive coastal resources.

However, the project as designed has the potential to result in adverse impacts to ESHA on site, given that it is located underneath the tree canopy of sensitive oak woodland habitat.

Therefore, in order to ensure that new development is consistent with the above referenced resource protection policies of both the Coastal Act and LUP, the Commission finds that **Special Condition One (1)** is necessary to ensure that temporary construction impacts to oak woodland habitat is minimized. Specifically, **Special Condition One (1)** requires that prior to issuance of the permit, the applicant shall submit, for the review and approval of the Executive Director, a detailed Oak Woodland Habitat Restoration Plan and Monitoring Program, prepared by a biologist or environmental resource specialist with qualifications acceptable to the Executive Director, for all areas of the project site temporarily disturbed by grading and

construction activities. Within 90 days of completion of the project, the applicant shall commence implementation of the approved riparian habitat restoration and mitigation plan. The Executive Director may grant additional time for good cause.

The Restoration Plan required pursuant to **Special Condition One (1)** shall provide for the mitigation for potential impacts to oak woodland habitat by replanting the understory areas that have been either temporarily disturbed or removed due to construction activities. These areas shall be planted with appropriate native plant species in the same location appropriate for an oak woodland area. All invasive and non-native plant species shall be removed from the project area and shall be revegetated with native plant species appropriate to an oak woodland habitat area. In addition, **Special Condition One (1)** also requires the applicant implement a five year monitoring program to ensure the success of the replanting.

Further, the project footprint is located in proximity to the trunks of three large coast live oak trees, representing a significant encroachment into the dripline of these tree canopies. Given the importance of oak woodlands and individual oak trees, even those that have been disturbed or fragmented by development, the Commission has consistently required, through past permit actions, that new development avoid the removal of oak trees, unless there is no feasible alternative for siting or designing the development. In this case, no oak trees will be removed. However, given the sensitivity of oak trees to disturbance or encroachment of development into the root zone, the Commission has required that encroachments within the protected zone (5 feet beyond the dripline, or 15 feet from the trunk, whichever is greater) be avoided unless there is no feasible alternative for the siting of development. The Commission has determined that in cases where the significant encroachment within the protected zone of oak trees is unavoidable, the planting of replacement trees is necessary as mitigation, at a ratio of at least ten seedlings for every tree impacted. If there is suitable area on the project site, replacement trees should be provided on-site. The Commission has found, through permit actions, based on the recommendations of resource specialists studying oak restoration that oak trees are most successfully established when planted as acorns collected in the local area or seedlings grown from such acorns. Many factors, over the life of the restoration, can result in the death of the replacement trees. In order to ensure that adequate replacement is eventually reached, it is necessary to provide a replacement ratio of at least ten replacement trees for every tree removed or impacted to account for the mortality of some of the replacement trees.

Therefore, in order to mitigate the impacts from the direct encroachment underneath the canopies of three oak trees, a total of 30 replacement trees must be planted. In order to provide this mitigation, the Commission finds it necessary to require the applicant to plant 30 replacement trees, as detailed in **Special Condition Two (2)**. **Special Condition 2** requires the applicant to plant at least 30 replacement seedlings, less than one year old, grown from acorns collected in the selected off-site planting area. Typically, the Commission will require such mitigation to be carried out on the project site, if suitable habitat exists therein. Since on-site mitigation is not feasible given the site's constrained topographic conditions and limited terms of access by the County of

Los Angeles, all replacement seedlings must be planted in a suitable location off-site that is restricted from development or is public parkland. An appropriate off-site mitigation area shall be identified within the Topanga Canyon watershed. **Special Condition 2** also requires the applicant to submit an off-site oak tree replacement planting program, which specifies replacement tree locations, tree or seedling size planting specifications, and a ten-year monitoring program to ensure that the replacement planting program is successful. The applicant may coordinate with the Mountains Restoration Trust, or other entity acceptable to the Executive Director, to prepare and implement the off-site oak tree replacement planting program. The applicant shall commence implementation of the approved off-site oak tree replacement planting program concurrently with the commencement of construction on the project site.

3. Water Quality

Section 30231 of the Coastal Act mandates the maintenance and protection of the biological productivity and quality of coastal waters and streams. The project area is situated several hundred feet upstream of the Topanga Canyon Creek channel that is considered to be ESHA and the potential exists for impacts to the water quality of the creek, particularly from erosion of sediment from the site. The sensitive habitats found in the stream and downstream of the project site could be adversely impacted by the proposed project through the introduction of excavated materials, chemicals, debris or sediment into the stream. Additionally, petroleum hydrocarbons including oil and grease from vehicles could be introduced to the stream through runoff. The discharge of these pollutants to coastal waters can cause cumulative impacts such as: eutrophication and anoxic conditions resulting in fish kills and diseases and the alteration of aquatic habitat including adverse changes to species composition and size; excess nutrients causing algae blooms and sedimentation increasing turbidity, which both reduce the penetration of sunlight needed by aquatic vegetation which provides food and cover for aquatic species; disruptions to the reproductive cycle of aquatic species; acute and sublethal toxicity in marine organisms leading to adverse changes in reproduction and feeding behavior; and human diseases such as hepatitis and dysentery. These impacts reduce the biological productivity and the quality of coastal waters, streams, wetlands, estuaries, and lakes and reduce optimum populations of marine organisms and have adverse impacts on human health.

To protect water quality adjacent to the project site during and after project construction, **Special Condition Six (6)** has been included to require the applicant to implement best management practices to minimize erosion and protect water quality. These measures include minimizing grading during the rainy season and implementing sediment control measures such as sand bag barriers, silt fencing, or geofabric covers on all cut or fill slopes, should construction activities occur during rainy season. **Special Condition 6** also requires that erosion measures be installed at the project site prior to or concurrent with the initial grading operations and maintained throughout the development process to minimize erosion and sediment from runoff waters during construction and requires

temporary erosion control measures should construction activities cease for more than 30 days.

Additionally, the project includes an excess of cut grading. These materials, if not properly managed and disposed, could have downstream water quality impacts. Excavated materials that are placed in stockpiles are subject to increased erosion. In order to ensure that excavated material will be limited to the designated temporary stockpile areas and be properly removed and disposed in a timely manner, **Special Condition Five (5)** also requires the applicant to properly remove all debris and excavated material from the site to a location that has authorization to receive such debris or fill.

The Commission finds that the proposed project, only as conditioned, will serve to maintain the quality of coastal waters to the extent possible and to minimize impacts to environmentally sensitive habitat area, while still allowing the repair project to proceed. It is thus as consistent as feasible with Sections 30231 and 30240 of the Coastal Act.

C. HAZARDS AND GEOLOGIC STABILITY

Coastal Act Section 30253 states in part:

New development shall:

- (1) Minimize risks to life and property in areas of high geologic, flood, and fire hazard.***
- (2) Assure stability and structural integrity, and neither create nor contribute significantly to erosion, geologic instability, or destruction of the site or surrounding area or in any way require the construction of protective devices that would substantially alter natural landforms along bluffs and cliffs.***

The proposed development is located in the Santa Monica Mountains, an area which is generally considered to be subject to an unusually high amount of natural hazards. Geologic hazards common to the Santa Monica Mountains include landslides, erosion, and flooding. In addition, fire is an inherent threat to the indigenous chaparral community of the coastal mountains. Wild fires often denude hillsides in the Santa Monica Mountains of all existing vegetation, thereby contributing to an increased potential for erosion and landslides on property.

The project includes reconstruction of an existing developed roadway and the construction of a new soldier pile retaining wall; after-the-fact authorization for the construction of a soldier pile retaining wall and reconstruction of the roadway along Grandview Drive; and revegetation of the temporary construction easement with native plant species after construction of the retaining wall has been completed.

The County Geotechnical and Materials Engineering Division has submitted an engineering memo for the proposed project, which indicates that the excessive runoff during the January 2005 storm season caused the road and hillside to fail along this section of Grandview Drive. The report noted that the large cracks in the roadway are due to "soil slumping caused by intense rainfall and ensuing saturation of the slope" and that "without construction of the soldier pile retaining wall the un-braced slope is susceptible to failure in the next occurrence of intense rainfall. Failure of the slope will render the roadway impassable and pose a public safety hazard." The purpose of the proposed project is to maintain the public's ability to use Grandview Drive for vehicular access and provide for emergency services/access to the residences and communities along this roadway.

The Commission notes that the proposed development, although necessary to remediate a hazardous eroding slope condition, will still not eliminate the potential for erosion of the steep slope on the subject site. The Commission finds that minimization of site erosion will add to the stability of the site. Erosion can best be minimized by requiring the applicant to plant all disturbed areas of the site with native plants compatible with the surrounding chaparral habitat. Further, in past permit actions, the Commission has found that invasive and non-native plant species are typically characterized as having a shallow root structure in comparison with their high surface/foilage weight and/or require a greater amount of irrigation and maintenance than native vegetation. The Commission notes that non-native and invasive plant species with high surface/foilage weight and shallow root structures do not serve to stabilize steep slopes, such as the slopes on the subject site, and that such vegetation results in potential adverse effects to the geologic stability of the project site. In comparison, the Commission finds that native plant species are typically characterized not only by a well-developed and extensive root structure in comparison to their surface/foilage weight but also by their low irrigation and maintenance requirements. Therefore, in order to ensure the stability and geotechnical safety of the site, **Special Condition One (1)** specifically requires that all proposed disturbed areas on subject site be stabilized with native vegetation appropriate for the oak woodland habitat.

The proposed project, as conditioned to ensure that the disturbed slopes on site are revegetated with native vegetation, has been designed to ensure slope stability on site to the maximum extent feasible. However, the Coastal Act recognizes that certain development projects located in geologically hazardous areas, such as the subject site, still involve the taking of some risk. Coastal Act policies require the Commission to establish the appropriate degree of risk acceptable for the proposed development and to determine who should assume the risk. When development in areas of identified hazards is proposed, the Commission considers the hazard associated with the project site and the potential cost to the public, as well as the individual's right to use his property. As such, the Commission finds that due to the foreseen possibility of erosion, landslide, and slope failure, the applicant shall assume these risks as a condition of approval. Therefore, **Special Condition Three (3)** requires the applicant to waive any claim of liability against the Commission for damage to life or property which may occur as a result of the permitted development. The applicant's assumption of risk, will show

that the applicant is aware of and appreciates the nature of the hazards which exist on the site, and which may adversely affect the stability or safety of the proposed development.

Therefore, for the reasons discussed above, the Commission finds that the proposed project, as conditioned, is consistent with Section 30253 of the Coastal Act.

D. VISUAL RESOURCES

Section 30251 of the Coastal Act states that:

The scenic and visual qualities of coastal areas shall be considered and protected as a resource of public importance. Permitted development shall be sited and designed to protect views to and along the ocean and scenic coastal areas, to minimize the alteration of natural land forms, to be visually compatible with the character of surrounding areas, and, where feasible, to restore and enhance visual quality in visually degraded areas. New development in highly scenic areas such as those designated in the California Coastline Preservation and Recreation Plan prepared by the Department of Parks and Recreation and by local government shall be subordinated to the character of its setting.

The project includes reconstruction of an existing developed roadway and the construction of a 71 linear-ft. soldier pile retaining wall; after-the-fact authorization for the construction of a 90 linear-ft. soldier pile retaining wall and reconstruction of the roadway along Grandview Drive; and revegetation of the temporary construction easement with native plant species after construction of the retaining wall has been completed. The existing 90 ft. wall was constructed in 1997 by Los Angeles County Public Works Department to restore the road after the 1995 storms caused slope failure below the road and damaged the roadway. The new wall would be a continuation of the south end of the existing unpermitted wall, and upon project completion, would appear as one integrated design.

The soldier pile retaining walls can be characterized as combination barrier railings, consisting of a 2-ft. 3-inch vehicular concrete barrier with a 2-ft. 3-inch bicycle rail on top of the concrete barrier (retaining wall). Approximately 4 ft. 6 inches would be exposed above grade and visible by the public along the roadway. The final wall would be visible by the public from immediate vantage points along the Grandview Drive roadway. The exterior of the wall (opposite side from the roadway) would be visible from the adjacent private development.

The project includes the removal of the existing 5 ft. high chain link fence on top of the concrete wall and subsequent replacement with the updated handrail design, thereby retrofitting the existing wall to meet the proposed maximum 4 ft. 6 inches height above the roadway. The accepted minimum guardrail height for pedestrians and wheelchairs is 42 inches above the pathway or sidewalk surface. (Caltrans Memo to Bridge Designers, Section 13-1). On bikeways, in addition to a handrail, most guidelines specify that an

additional rail, a higher guardrail, be provided to prevent toppling, since cyclists have a higher center of gravity than pedestrians. Generally, the recommended bicycle guardrail height is 4 ft. 6 inches (Caltrans Highway Design Manual, page 200-40). Caltrans allows that a lower rail may be used if a curbed sidewalk separates the bikeway from the railing (Caltrans Highway Design Manual, page 200-41). Pedestrian and/or bicycle railings are required for safety reasons where the vertical fall is 4 feet or more adjacent to a retaining wall. In this case, the vertical drop would be more than 4 feet on the outboard slope side of the wall, and therefore a railing is required.

The proposed concrete wall will serve as a barrier for vehicles, while the proposed handrail is necessary to ensure protection of pedestrians and bicyclists. The applicant has worked with Commission staff to provide design alternatives that would meet the safety design requirements while at the same time reduce the visual intrusion of the barrier, given the rural community character of the subject site. The potential alternative designs included wrought-iron type fence, tubular handrail design (i.e., proposed design), or chain link fence. An alternative concrete rail design was not visually permeable and therefore was not considered. Additionally, the width of the roadway was too constrained to support a separate walkway/bikeway that would allow separation of the vehicular & bicycle traffic from pedestrian, and thus, the potential for a lower overall rail design was not feasible given these circumstances. Staff reviewed the alternatives and determined that the handrail, if modified to blend with the natural surroundings, would be the alternative most in keeping with the rural community.

The Commission finds that the soldier pile retaining wall, road reconstruction, and associated grading would serve to increase the structural stability of the roadway on the subject site and ensure public safety. However, the wall will be visible along Grandview Drive (a public roadway), and the wall would be more urban in appearance and less consistent with the rural nature of the area surrounding the project site than previously existed. Therefore, to minimize adverse effects to public views resulting from the proposed development, the Commission finds that the surface of the proposed soldier pile retaining wall and handrail barrier must be designed to include, blend, or mimic, the color and texture of native materials and appearance of the natural environment (such as the appearance of rock facing), pursuant to **Special Condition Four (4)**.

For the reasons discussed above, the Commission finds that the proposed development, as conditioned, protects the scenic and visual qualities of the area and is consistent with Section 30251 of the Coastal Act.

E. UNPERMITTED DEVELOPMENT

Development has occurred on Grandview Drive adjacent to the subject site without the required coastal development permit, including but not limited to construction of a 90 linear foot soldier pile retaining wall, 2 ft. 3 inches in height with a five-foot chain link fence on top, reconstruction of approximately 1,210 sq. ft. of Grandview Drive roadway, and 89 cu. yds. (56 cu. yds cut, and 33 cu. yds. fill) of grading. The wall was constructed in 1997 by Los Angeles County Public Works Department to restore the road after the

1995 storms caused slope failure below the road and damaged the roadway. This application includes a request for after-the-fact authorization for the above referenced unpermitted development. The area where the existing wall is located is directly adjacent to residential development including building constructed prior to the Coastal Act, and although located under the canopy of oak trees that are situated on the other side of Grandview Drive, there are no oak trees directly located along this stretch of the project. In order to ensure that the unpermitted development component of this application is resolved in a timely manner, the Commission finds it necessary to require, through **Special Condition Seven (7)**, the applicant to fulfill all of the Special Conditions as a prerequisite to the issuance of this permit within 180 days of Commission action.

F. LOCAL COASTAL PROGRAM

Section 30604 of the Coastal Act states:

a) Prior to certification of the local coastal program, a coastal development permit shall be issued if the issuing agency, or the commission on appeal, finds that the proposed development is in conformity with the provisions of Chapter 3 (commencing with Section 30200) of this division and that the permitted development will not prejudice the ability of the local government to prepare a local program that is in conformity with the provisions of Chapter 3 (commencing with Section 30200).

Section 30604(a) of the Coastal Act provides that the Commission shall issue a Coastal Development Permit only if the project will not prejudice the ability of the local government having jurisdiction to prepare a Local Coastal Program that conforms with the Chapter 3 policies of the Coastal Act. The preceding sections provide findings that the proposed project will be in conformity with the provisions of Chapter 3 if certain conditions are incorporated into the project and are accepted by the applicant. As conditioned, the proposed development will not create adverse impacts and is found to be consistent with the applicable policies contained in Chapter 3. Therefore, the Commission finds that approval of the proposed development, as conditioned, will not prejudice the County of Los Angeles' ability to prepare a Local Coastal Program for this area which is also consistent with the policies of Chapter 3 of the Coastal Act, as required by Section 30604(a).

G. CEQA

Section 13096(a) of the Commission's administrative regulations requires Commission approval of a Coastal Development Permit application to be supported by a finding showing the application, as conditioned by any conditions of approval, to be consistent with any applicable requirements of the California Environmental Quality Act (CEQA). Section 21080.5(d)(2)(A) of CEQA prohibits a proposed development from being approved if there are feasible alternatives or feasible mitigation measures available

which would substantially lessen any significant adverse effect that the activity may have on the environment.

The County of Los Angeles found that the proposed project was statutorily exempt pursuant to Section 21080 (b) (3) of the California Environmental Quality.

The Commission incorporates its findings on Coastal Act consistency at this point as if set forth in full. These findings address and respond to all public comments regarding potential significant adverse environmental effects of the project that were received prior to preparation of the staff report. As discussed above, the proposed development is authorized by the Act and, as conditioned, is as consistent with the policies of the Coastal Act as is possible. Feasible mitigation measures which will minimize all adverse environmental effects have been required as special conditions. As conditioned, there are no feasible alternatives or feasible mitigation measures available, beyond those required, which would substantially lessen any significant adverse impact that the activity may have on the environment. Therefore, the Commission finds that the proposed project, as conditioned to mitigate the identified impacts, can be found to be consistent with the requirements of the Coastal Act to conform to CEQA.

GRANDVIEW DRIVE-60 FEET NORTH OF FALLS DRIVE LOCATION MAP





Exhibit 1
4-06-153
Vicinity Map

... TO PLANS



PLAN
1" = 30'

CURVE DATA			
Δ	T	L	
14° 52' 35"	25.46	50.83	
30° 26' 30"	28.57	55.79	
47° 51' 30"	24.41	45.94	
45° 18' 00"	10.68	20.19	

Exhibit 2a
4-06-153
Site Plan
New 71 linear-ft Wall

Note:
Removal of existing guard rail, dke, and chainlink fence for wall construction are not shown for clarity. See Special Provisions.

BENCH MARK

MAN BU 1980
FD MARIAL & SM WISHER ± 208 FT WLY OF
INT GRANDVIEW DR & FALLS DR & 3.5 FT WLY
FROM WALL & ELY SIDE OF GRANDVIEW DR



COUNTY OF LOS ANGELES DEPARTMENT OF PUBLIC WORKS




GRANDVIEW DRIVE
60' N/O FALLS DRIVE

PROJECT ID NO. RDC0014683
PLAN

THE CONTRACTOR SHALL VERIFY ALL CONTROLLING FIELD DIMENSIONS BEFORE ORDERING OR FABRICATING ANY MATERIAL.


DATE	BY	REVISIONS

BR NO. 3998 PCA RB3135062 DMC P8519338 SHEET 3 OF 6



EXIST. WALL ELEVATION

1" = 10' - 0"




NEW WALL ELEVATION

1" = 10'

Exhibit 3
4-06-153
Wall Plans

See Revised Railing Plan
(Exhibit 4)

Q Pile & Wall LOL



NOTE:

1. All exposed structural steel shall be painted with zinc-based paint. See Special Provisions.
2. Edge of pavement elevation may be adjusted to smooth roadway surface with an approval from the Engineer.

SECTION A-A

1/4" = 1'-0"

THE CONTRACTOR SHALL VERIFY ALL CONTROLLING FIELD DIMENSIONS BEFORE ORDERING OR FABRICATING ANY MATERIAL.

REVIEWED BY DATE 12/1/01

PROJECT FILE NAME R0004683-BRD

DESIGNED BY T. SANDEFUR

CHECKED BY E.F. RODRIGUEZ

DRAWN BY L. GRINSHTEYN





Exhibit 4
4-06-153
Railing Plan



SECTION A-A

CALIFORNIA COASTAL COMMISSION

45 FREMONT, SUITE 2000
SAN FRANCISCO, CA 94105-2219
VOICE AND TDD (415) 904-5200
FAX (415) 904-5400



M E M O R A N D U M

FROM: John Dixon, Ph.D.
Ecologist / Wetland Coordinator

TO: Ventura Staff

SUBJECT: Designation of ESHA in the Santa Monica Mountains

DATE: March 25, 2003

In the context of the Malibu LCP, the Commission found that the Mediterranean Ecosystem in the Santa Mountains is rare, and especially valuable because of its relatively pristine character, physical complexity, and resultant biological diversity. Therefore, areas of undeveloped native habitat in the Santa Monica Mountains that are large and relatively unfragmented may meet the definition of ESHA by virtue of their valuable roles in that ecosystem, regardless of their relative rarity throughout the state. This is the only place in the coastal zone where the Commission has recognized chaparral as meeting the definition of ESHA. The scientific background presented herein for ESHA analysis in the Santa Monica Mountains is adapted from the Revised Findings for the Malibu LCP that the Commission adopted on February 6, 2003.

For habitats in the Santa Monica Mountains, particularly coastal sage scrub and chaparral, there are three site-specific tests to determine whether an area is ESHA because of its especially valuable role in the ecosystem. First, is the habitat properly identified, for example as coastal sage scrub or chaparral? The requisite information for this test generally should be provided by a site-specific biological assessment. Second, is the habitat largely undeveloped and otherwise relatively pristine? Third, is the habitat part of a large, contiguous block of relatively pristine native vegetation? This should be documented with an aerial photograph from our mapping unit (with the site delineated) and should be attached as an exhibit to the staff report. For those habitats that are absolutely rare or that support individual rare species, it is not necessary to find that they are relatively pristine, and are neither isolated nor fragmented.

**Designation of Environmentally Sensitive Habitat in the
Santa Monica Mountains**

The Coastal Act provides a definition of “environmentally sensitive area” as: “Any area in which plant or animal life or their habitats are either rare or especially valuable because of their special nature or role in an ecosystem and which could be easily disturbed or degraded by human activities and developments” (Section 30107.5).

Exhibit 5 4-06-153 Dr. Dixon ESHA Memo
--

There are three important elements to the definition of ESHA. First, a geographic area can be designated ESHA either because of the presence of individual species of plants or animals or because of the presence of a particular habitat. Second, in order for an area to be designated as ESHA, the species or habitat must be either rare or it must be especially valuable. Finally, the area must be easily disturbed or degraded by human activities.

The first test of ESHA is whether a habitat or species is rare. Rarity can take several forms, each of which is important. Within the Santa Monica Mountains, rare species and habitats often fall within one of two common categories. Many rare species or habitats are globally rare, but locally abundant. They have suffered severe historical declines in overall abundance and currently are reduced to a small fraction of their original range, but where present may occur in relatively large numbers or cover large local areas. This is probably the most common form of rarity for both species and habitats in California and is characteristic of coastal sage scrub, for example. Some other habitats are geographically widespread, but occur everywhere in low abundance. California's native perennial grasslands fall within this category.

A second test for ESHA is whether a habitat or species is especially valuable. Areas may be valuable because of their "special nature," such as being an unusually pristine example of a habitat type, containing an unusual mix of species, supporting species at the edge of their range, or containing species with extreme variation. For example, reproducing populations of valley oaks are not only increasingly rare, but their southernmost occurrence is in the Santa Monica Mountains. Generally, however, habitats or species are considered valuable because of their special "role in the ecosystem." For example, many areas within the Santa Monica Mountains may meet this test because they provide habitat for endangered species, protect water quality, provide essential corridors linking one sensitive habitat to another, or provide critical ecological linkages such as the provision of pollinators or crucial trophic connections. Of course, all species play a role in their ecosystem that is arguably "special." However, the Coastal Act requires that this role be "especially valuable." This test is met for relatively pristine areas that are integral parts of the Santa Monica Mountains Mediterranean ecosystem because of the demonstrably rare and extraordinarily special nature of that ecosystem as detailed below.

Finally, ESHAs are those areas that could be easily disturbed or degraded by human activities and developments. Within the Santa Monica Mountains, as in most areas of southern California affected by urbanization, all natural habitats are in grave danger of direct loss or significant degradation as a result of many factors related to anthropogenic changes.

Ecosystem Context of the Habitats of the Santa Monica Mountains

The Santa Monica Mountains comprise the largest, most pristine, and ecologically complex example of a Mediterranean ecosystem in coastal southern California.

California's coastal sage scrub, chaparral, oak woodlands, and associated riparian areas have analogues in just a few areas of the world with similar climate. Mediterranean ecosystems with their wet winters and warm dry summers are only found in five localities (the Mediterranean coast, California, Chile, South Africa, and south and southwest Australia). Throughout the world, this ecosystem with its specially adapted vegetation and wildlife has suffered severe loss and degradation from human development. Worldwide, only 18 percent of the Mediterranean community type remains undisturbed¹. However, within the Santa Monica Mountains, this ecosystem is remarkably intact despite the fact that it is closely surrounded by some 17 million people. For example, the 150,000 acres of the Santa Monica Mountains National Recreation Area, which encompasses most of the Santa Monica Mountains, was estimated to be 90 percent free of development in 2000². Therefore, this relatively pristine area is both large and mostly unfragmented, which fulfills a fundamental tenet of conservation biology³. The need for large contiguous areas of natural habitat in order to maintain critical ecological processes has been emphasized by many conservation biologists⁴.

In addition to being a large single expanse of land, the Santa Monica Mountains ecosystem is still connected, albeit somewhat tenuously, to adjacent, more inland ecosystems⁵. Connectivity among habitats within an ecosystem and connectivity among ecosystems is very important for the preservation of species and ecosystem integrity. In a recent statewide report, the California Resources Agency⁶ identified wildlife corridors and habitat connectivity as the top conservation priority. In a letter to governor Gray Davis, sixty leading environmental scientists have endorsed the

¹ National Park Service. 2000. Draft general management plan & environmental impact statement. Santa Monica Mountains National Recreation Area – California.

² Ibid.

³ Harris, L. D. 1988. Edge effects and conservation of biotic diversity. *Conserv. Biol.* 330-332. Soule, M. E, D. T. Bolger, A. C. Alberts, J. Wright, M. Sorice and S. Hill. 1988. Reconstructed dynamics of rapid extinctions of chaparral-requiring birds in urban habitat islands. *Conserv. Biol.* 2: 75-92. Yahner, R. H. 1988. Changes in wildlife communities near edges. *Conserv. Biol.* 2:333-339. Murphy, D. D. 1989. Conservation and confusion: Wrong species, wrong scale, wrong conclusions. *Conservation Biol.* 3:82-84.

⁴ Crooks, K. 2000. Mammalian carnivores as target species for conservation in Southern California. p. 105-112 in: Keeley, J. E., M. Baer-Keeley and C. J. Fotheringham (eds), 2nd Interface Between Ecology and Land Development in California, U.S. Geological Survey Open-File Report 00-62. Sauvajot, R. M., E. C. York, T. K. Fuller, H. Sharon Kim, D. A. Kamradt and R. K. Wayne. 2000. Distribution and status of carnivores in the Santa Monica Mountains, California: Preliminary results from radio telemetry and remote camera surveys. p 113-123 in: Keeley, J. E., M. Baer-Keeley and C. J. Fotheringham (eds), 2nd Interface Between Ecology and Land Development in California, U.S. Geological Survey Open-File Report 00-62. Beier, P. and R. F. Noss. 1998. Do habitat corridors provide connectivity? *Conserv. Biol.* 12:1241-1252. Beier, P. 1996. Metapopulation models, tenacious tracking and cougar conservation. In: *Metapopulations and Wildlife Conservation*, ed. D. R. McCullough. Island Press, Covelo, California, 429p.

⁵ The SMM area is linked to larger natural inland areas to the north through two narrow corridors: 1) the Conejo Grade connection at the west end of the Mountains and 2) the Simi Hills connection in the central region of the SMM (from Malibu Creek State Park to the Santa Susanna Mountains).

⁶ California Resources Agency. 2001. Missing Linkages: Restoring Connectivity to the California Landscape. California Wilderness Coalition, Calif. Dept of Parks & Recreation, USGS, San Diego Zoo and The Nature Conservancy. Available at: <http://www.calwild.org/pubs/reports/linkages/index.htm>

conclusions of that report⁷. The chief of natural resources at the California Department of Parks and Recreation has identified the Santa Monica Mountains as an area where maintaining connectivity is particularly important⁸.

The species most directly affected by large scale connectivity are those that require large areas or a variety of habitats, e.g., gray fox, cougar, bobcat, badger, steelhead trout, and mule deer⁹. Large terrestrial predators are particularly good indicators of habitat connectivity and of the general health of the ecosystem¹⁰. Recent studies show that the mountain lion, or cougar, is the most sensitive indicator species of habitat fragmentation, followed by the spotted skunk and the bobcat¹¹. Sightings of cougars in both inland and coastal areas of the Santa Monica Mountains¹² demonstrate their continued presence. Like the “canary in the mineshaft,” an indicator species like this is good evidence that habitat connectivity and large scale ecological function remains in the Santa Monica Mountains ecosystem.

The habitat integrity and connectivity that is still evident within the Santa Monica Mountains is extremely important to maintain, because both theory and experiments over 75 years in ecology confirm that large spatially connected habitats tend to be more stable and have less frequent extinctions than habitats without extended spatial structure¹³. Beyond simply destabilizing the ecosystem, fragmentation and disturbance

⁷ Letters received and included in the September 2002 staff report for the Malibu LCP.

⁸ Schoch, D. 2001. Survey lists 300 pathways as vital to state wildlife. Los Angeles Times. August 7, 2001.

⁹ Martin, G. 2001. Linking habitat areas called vital for survival of state's wildlife Scientists map main migration corridors. San Francisco Chronicle, August 7, 2001.

¹⁰ Noss, R. F., H. B. Quigley, M. G. Hornocker, T. Merrill and P. C. Paquet. 1996. Conservation biology and carnivore conservation in the Rocky Mountains. *Conserv. Biol.* 10: 949-963. Noss, R. F. 1995. Maintaining ecological integrity in representative reserve networks. World Wildlife Fund Canada.

¹¹ Sauvajot, R. M., E. C. York, T. K. Fuller, H. Sharon Kim, D. A. Kamradt and R. K. Wayne. 2000. Distribution and status of carnivores in the Santa Monica Mountains, California: Preliminary results from radio telemetry and remote camera surveys. p 113-123 in: Keeley, J. E., M. Baer-Keeley and C. J. Fotheringham (eds), 2nd Interface Between Ecology and Land Development in California, U.S. Geological Survey Open-File Report 00-62. Beier, P. 1996. Metapopulation models, tenacious tracking and cougar conservation. In: *Metapopulations and Wildlife Conservation*, ed. D. R. McCullough. Island Press, Covelo, California, 429p.

¹² Recent sightings of mountain lions include: Temescal Canyon (pers. com., Peter Brown, Facilities Manager, Calvary Church), Topanga Canyon (pers. com., Marti Witter, NPS), Encinal and Trancas Canyons (pers. com., Pat Healy), Stump Ranch Research Center (pers. com., Dr. Robert Wayne, Dept. of Biology, UCLA). In May of 2002, the NPS *photographed* a mountain lion at a trip camera on the Back Bone Trail near Castro Crest – Seth Riley, Eric York and Dr. Ray Sauvajot, National Park Service, SMMNRA.

¹³ Gause, G. F. 1934. The struggle for existence. Baltimore, William and Wilkins 163 p. (also reprinted by Hafner, N.Y. 1964). Gause, G. F., N. P. Smaragdova and A. A. Witt. 1936. Further studies of interaction between predators and their prey. *J. Anim. Ecol.* 5:1-18. Huffaker, C. B. 1958. Experimental studies on predation: dispersion factors and predator-prey oscillations. *Hilgardia* 27:343-383. Luckinbill, L. S. 1973. Coexistence in laboratory populations of *Paramecium aurelia* and its predator *Didinium nasutum*. *Ecology* 54:1320-1327. Allen, J. C., C. C. Brewster and D. H. Slone. 2001. Spatially explicit ecological models: A spatial convolution approach. *Chaos, Solitons and Fractals*. 12:333-347.

can even cause unexpected and irreversible changes to new and completely different kinds of ecosystems (habitat conversion)¹⁴.

As a result of the pristine nature of large areas of the Santa Monica Mountains and the existence of large, unfragmented and interconnected blocks of habitat, this ecosystem continues to support an extremely diverse flora and fauna. The observed diversity is probably a function of the diversity of physical habitats. The Santa Monica Mountains have the greatest geological diversity of all major mountain ranges within the transverse range province. According to the National Park Service, the Santa Monica Mountains contain 40 separate watersheds and over 170 major streams with 49 coastal outlets¹⁵. These streams are somewhat unique along the California coast because of their topographic setting. As a "transverse" range, the Santa Monica Mountains are oriented in an east-west direction. As a result, the south-facing riparian habitats have more variable sun exposure than the east-west riparian corridors of other sections of the coast. This creates a more diverse moisture environment and contributes to the higher biodiversity of the region. The many different physical habitats of the Santa Monica Mountains support at least 17 native vegetation types¹⁶ including the following habitats considered sensitive by the California Department of Fish and Game: native perennial grassland, coastal sage scrub, red-shank chaparral, valley oak woodland, walnut woodland, southern willow scrub, southern cottonwood-willow riparian forest, sycamore-alder woodland, oak riparian forest, coastal salt marsh, and freshwater marsh. Over 400 species of birds, 35 species of reptiles and amphibians, and more than 40 species of mammals have been documented in this diverse ecosystem. More than 80 sensitive species of plants and animals (listed, proposed for listing, or species of concern) are known to occur or have the potential to occur within the Santa Monica Mountains Mediterranean ecosystem.

The Santa Monica Mountains are also important in a larger regional context. Several recent studies have concluded that the area of southern California that includes the Santa Monica Mountains is among the most sensitive in the world in terms of the number of rare endemic species, endangered species and habitat loss. These studies have designated the area to be a local hot-spot of endangerment in need of special protection¹⁷.

Therefore, the Commission finds that the Santa Monica Mountains ecosystem is itself rare and especially valuable because of its special nature as the largest, most pristine,

¹⁴ Scheffer, M., S. Carpenter, J. A. Foley, C. Folke and B. Walker. 2001. Catastrophic shifts in ecosystems. *Nature* 413:591-596.

¹⁵ NPS. 2000. op.cit.

¹⁶ From the NPS report (2000 op. cit.) that is based on the older Holland system of subjective classification. The data-driven system of Sawyer and Keeler-Wolf results in a much larger number of distinct "alliances" or vegetation types.

¹⁷ Myers, N. 1990. The biodiversity challenge: Expanded hot-spots analysis. *Environmentalist* 10:243-256. Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca and J. A. Kent. 2000. Biodiversity hot-spots for conservation priorities. *Nature* 403:853-858. Dobson, A. P., J. P. Rodriguez, W. M. Roberts and D. S. Wilcove. 1997. Geographic distribution of endangered species in the United States. *Science* 275:550-553.

physically complex, and biologically diverse example of a Mediterranean ecosystem in coastal southern California. The Commission further finds that because of the rare and special nature of the Santa Monica Mountains ecosystem, the ecosystem roles of substantially intact areas of the constituent plant communities discussed below are “especially valuable” under the Coastal Act.

Major Habitats within the Santa Monica Mountains

The most recent vegetation map that is available for the Santa Monica Mountains is the map that was produced for the National Park Service in the mid-1990s using 1993 satellite imagery supplemented with color and color infrared aerial imagery from 1984, 1988, and 1994 and field review¹⁸. The minimum mapping unit was 5 acres. For that map, the vegetation was mapped in very broad categories, generally following a vegetation classification scheme developed by Holland¹⁹. Because of the mapping methods used the degree of plant community complexity in the landscape is not represented. For example, the various types of “ceanothus chaparral” that have been documented were lumped under one vegetation type referred to as “northern mixed chaparral.” Dr. Todd Keeler-Wolf of the California Department of Fish and Game is currently conducting a more detailed, quantitative vegetation survey of the Santa Monica Mountains.

The National Park Service map can be used to characterize broadly the types of plant communities present. The main generic plant communities present in the Santa Monica Mountains²⁰ are: coastal sage scrub, chaparral, riparian woodland, coast live oak woodland, and grasslands.

Riparian Woodland

Some 49 streams connect inland areas with the coast, and there are many smaller drainages as well, many of which are “blue line.” Riparian woodlands occur along both perennial and intermittent streams in nutrient-rich soils. Partly because of its multi-layered vegetation, the riparian community contains the greatest overall biodiversity of all the plant communities in the area²¹. At least four types of riparian communities are discernable in the Santa Monica Mountains: walnut riparian areas, mulefat-dominated riparian areas, willow riparian areas and sycamore riparian woodlands. Of these, the

¹⁸ Franklin, J. 1997. Forest Service Southern California Mapping Project, Santa Monica Mountains National Recreation Area, Task 11 Description and Results, Final Report. June 13, 1997, Dept. of Geography, San Diego State University, USFS Contract No. 53-91S8-3-TM45.

¹⁹ Holland R. F. 1986. Preliminary Descriptions of the Terrestrial Natural Communities of California. State of California, The Resources Agency, Dept. of Fish and Game, Natural Heritage Division, Sacramento, CA. 95814.

²⁰ National Park Service. 2000. Draft: General Management Plan & Environmental Impact Statement, Santa Monica Mountains National Recreation Area, US Dept. of Interior, National Park Service, December 2000. (Fig. 11 in this document.)

²¹ Ibid.

sycamore riparian woodland is the most diverse riparian community in the area. In these habitats, the dominant plant species include arroyo willow, California black walnut, sycamore, coast live oak, Mexican elderberry, California bay laurel, and mule fat. Wildlife species that have been observed in this community include least Bell's vireo (a State and federally listed species), American goldfinches, black phoebes, warbling vireos, bank swallows (State listed threatened species), song sparrows, belted kingfishers, raccoons, and California and Pacific tree frogs.

Riparian communities are the most species-rich to be found in the Santa Monica Mountains. Because of their multi-layered vegetation, available water supply, vegetative cover and adjacency to shrubland habitats, they are attractive to many native wildlife species, and provide essential functions in their lifecycles²². During the long dry summers in this Mediterranean climate, these communities are an essential refuge and oasis for much of the areas' wildlife.

Riparian habitats and their associated streams form important connecting links in the Santa Monica Mountains. These habitats connect all of the biological communities from the highest elevation chaparral to the sea with a unidirectional flowing water system, one function of which is to carry nutrients through the ecosystem to the benefit of many different species along the way.

The streams themselves provide refuge for sensitive species including: the coast range newt, the Pacific pond turtle, and the steelhead trout. The coast range newt and the Pacific pond turtle are California Species of Special Concern and are proposed for federal listing²³, and the steelhead trout is federally endangered. The health of the streams is dependent on the ecological functions provided by the associated riparian woodlands. These functions include the provision of large woody debris for habitat, shading that controls water temperature, and input of leaves that provide the foundation of the stream-based trophic structure.

The importance of the connectivity between riparian areas and adjacent habitats is illustrated by the Pacific pond turtle and the coast range newt, both of which are sensitive and both of which require this connectivity for their survival. The life history of the Pacific pond turtle demonstrates the importance of riparian areas and their associated watersheds for this species. These turtles require the stream habitat during the wet season. However, recent radio tracking work²⁴ has found that although the Pacific pond turtle spends the wet season in streams, it also requires upland habitat for refuge during the dry season. Thus, in coastal southern California, the Pacific pond turtle requires both streams and intact adjacent upland habitats such as coastal sage

²² Walter, Hartmut. Bird use of Mediterranean habitats in the Santa Monica Mountains, Coastal Commission Workshop on the Significance of Native Habitats in the Santa Monica Mountains. CCC Hearing, June 13, 2002, Queen Mary Hotel.

²³ USFWS. 1989. Endangered and threatened wildlife and plants; animal notice of review. Fed. Reg. 54:554-579. USFWS. 1993. Endangered and threatened wildlife and plants; notice of 1-year petition finding on the western pond turtle. Fed. Reg. 58:42717-42718.

²⁴ Rathbun, G.B., N.J. Scott and T.G. Murphy. 2002. Terrestrial habitat use by Pacific pond turtle in a Mediterranean climate. *Southwestern Naturalist*. (in Press).

scrub, woodlands or chaparral as part of their normal life cycle. The turtles spend about four months of the year in upland refuge sites located an average distance of 50 m (but up to 280 m) from the edge of the creek bed. Similarly, nesting sites where the females lay eggs are also located in upland habitats an average of 30 m (but up to 170 m) from the creek. Occasionally, these turtles move up to 2 miles across upland habitat²⁵. Like many species, the pond turtle requires both stream habitats and the upland habitats of the watershed to complete its normal annual cycle of behavior. Similarly, the coast range newt has been observed to travel hundreds of meters into upland habitat and spend about ten months of the year far from the riparian streambed²⁶. They return to the stream to breed in the wet season, and they are therefore another species that requires both riparian habitat and adjacent uplands for their survival.

Riparian habitats in California have suffered serious losses and such habitats in southern California are currently very rare and seriously threatened. In 1989, Faber estimated that 95-97% of riparian habitat in southern California was already lost²⁷. Writing at the same time as Faber, Bowler asserted that, "[t]here is no question that riparian habitat in southern California is endangered."²⁸ In the intervening 13 years, there have been continuing losses of the small amount of riparian woodlands that remain. Today these habitats are, along with native grasslands and wetlands, among the most threatened in California.

In addition to direct habitat loss, streams and riparian areas have been degraded by the effects of development. For example, the coast range newt, a California Species of Special Concern has suffered a variety of impacts from human-related disturbances²⁹. Human-caused increased fire frequency has resulted in increased sedimentation rates, which exacerbates the cannibalistic predation of adult newts on the larval stages.³⁰ In addition impacts from non-native species of crayfish and mosquito fish have also been documented. When these non-native predators are introduced, native prey organisms are exposed to new mortality pressures for which they are not adapted. Coast range newts that breed in the Santa Monica Mountain streams do not appear to have adaptations that permit co-occurrence with introduced mosquito fish and crayfish³¹. These introduced predators have eliminated the newts from streams where they previously occurred by both direct predation and suppression of breeding.

²⁵ Testimony by R. Dagit, Resource Conservation District of the Santa Monica Mountains at the CCC Habitat Workshop on June 13, 2002.

²⁶ Dr. Lee Kats, Pepperdine University, personal communication to Dr J. Allen, CCC.

²⁷ Faber, P.A., E. Keller, A. Sands and B.M. Massey. 1989. The ecology of riparian habitats of the southern California coastal region: a community profile. U.S. Fish and Wildlife Service Biological Report 85(7.27) 152pp.

²⁸ Bowler, P.A. 1989. Riparian woodland: An endangered habitat in southern California. Pp 80-97 in Schoenherr, A.A. (ed.) Endangered plant communities of southern California. Botanists Special Publication No. 3.

²⁹ Gamradt, S.C., L.B. Kats and C.B. Anzalone. 1997. Aggression by non-native crayfish deters breeding in California newts. *Conservation Biology* 11(3):793-796.

³⁰ Kerby, L.J., and L.B. Kats. 1998. Modified interactions between salamander life stages caused by wildfire-induced sedimentation. *Ecology* 79(2):740-745.

³¹ Gamradt, S.C. and L.B. Kats. 1996. Effect of introduced crayfish and mosquitofish on California newts. *Conservation Biology* 10(4):1155-1162.

Therefore, because of the essential role that riparian plant communities play in maintaining the biodiversity of the Santa Monica Mountains, because of the historical losses and current rarity of these habitats in southern California, and because of their extreme sensitivity to disturbance, the native riparian habitats in the Santa Monica Mountains meet the definition of ESHA under the Coastal Act.

Coastal Sage Scrub and Chaparral

Coastal sage scrub and chaparral are often lumped together as “shrublands” because of their roughly similar appearance and occurrence in similar and often adjacent physical habitats. In earlier literature, these vegetation associations were often called soft chaparral and hard chaparral, respectively. “Soft” and “hard” refers to differences in their foliage associated with different adaptations to summer drought. Coastal sage scrub is dominated by soft-leaved, generally low-growing aromatic shrubs that die back and drop their leaves in response to drought. Chaparral is dominated by taller, deeper-rooted evergreen shrubs with hard, waxy leaves that minimize water loss during drought.

The two vegetation types are often found interspersed with each other. Under some circumstances, coastal sage scrub may even be successional to chaparral, meaning that after disturbance, a site may first be covered by coastal sage scrub, which is then replaced with chaparral over long periods of time.³² The existing mosaic of coastal sage scrub and chaparral is the result of a dynamic process that is a function of fire history, recent climatic conditions, soil differences, slope, aspect and moisture regime, and the two habitats should not be thought of as completely separate and unrelated entities but as different phases of the same process³³. The spatial pattern of these vegetation stands at any given time thus depends on both local site conditions and on history (e.g., fire), and is influenced by both natural and human factors.

In lower elevation areas with high fire frequency, chaparral and coastal sage scrub may be in a state of flux, leading one researcher to describe the mix as a “coastal sage-chaparral subclimax.”³⁴ Several other researchers have noted the replacement of chaparral by coastal sage scrub, or coastal sage scrub by chaparral depending on fire history.³⁵ In transitional and other settings, the mosaic of chaparral and coastal sage

³² Cooper, W.S. 1922. The broad-sclerophyll vegetation of California. Carnegie Institution of Washington Publication 319. 124 pp.

³³ Longcore, T and C. Rich. 2002. Protection of environmentally sensitive habitat areas in proposed local coastal plan for the Santa Monica Mountains. The Urban Wildlands Group, Inc., P.O. Box 24020 Los Angeles, CA 90024. (See attached comment document in Appendix).

³⁴ Hanes, T.L. 1965. Ecological studies on two closely related chaparral shrubs in southern California. Ecological Monographs 41:27-52.

³⁵ Gray, K.L. 1983. Competition for light and dynamic boundary between chaparral and coastal sage scrub. Madrono 30(1):43-49. Zedler, P.H., C.R. Gautier and G.S. McMaster. 1983. Vegetation change in response to extreme events: The effect of a short interval between fires in California chaparral and coastal sage scrub. Ecology 64(4): 809-818.

scrub enriches the seasonal plant resource base and provides additional habitat variability and seasonality for the many species that inhabit the area.

Relationships Among Coastal Sage Scrub, Chaparral and Riparian Communities

Although the constituent communities of the Santa Monica Mountains Mediterranean ecosystem can be defined and distinguished based on species composition, growth habits, and the physical habitats they characteristically occupy, they are not independent entities ecologically. Many species of plants, such as black sage, and laurel sumac, occur in more than one plant community and many animals rely on the predictable mix of communities found in undisturbed Mediterranean ecosystems to sustain them through the seasons and during different portions of their life histories.

Strong evidence for the interconnectedness between chaparral, coastal scrub and other habitats is provided by “opportunistic foragers” (animals that follow the growth and flowering cycles across these habitats). Coastal scrub and chaparral flowering and growth cycles differ in a complimentary and sequential way that many animals have evolved to exploit. Whereas coastal sage scrub is shallow-rooted and responds quickly to seasonal rains, chaparral plants are typically deep-rooted having most of their flowering and growth later in the rainy season after the deeper soil layers have been saturated³⁶. New growth of chaparral evergreen shrubs takes place about four months later than coastal sage scrub plants and it continues later into the summer³⁷. For example, in coastal sage scrub, California sagebrush flowers and grows from August to February and coyote bush flowers from August to November³⁸. In contrast, chamise chaparral and bigpod ceanothus flower from April to June, buck brush ceanothus flowers from February to April, and hoaryleaf ceanothus flowers from March to April.

Many groups of animals exploit these seasonal differences in growth and blooming period. The opportunistic foraging insect community (e.g., honeybees, butterflies and moths) tends to follow these cycles of flowering and new growth, moving from coastal sage scrub in the early rainy season to chaparral in the spring³⁹. The insects in turn are followed by insectivorous birds such as the blue-gray gnatcatcher⁴⁰, bushtit, cactus wren, Bewick’s wren and California towhee. At night bats take over the role of daytime insectivores. At least 12 species of bats (all of which are considered sensitive) occur in

³⁶ DeSimone, S. 2000. California’s coastal sage scrub. *Fremontia* 23(4):3-8. Mooney, H.A. 1988. Southern coastal scrub. Chap. 13 in Barbour, M.G. and J. Majors; Eds. 1988. *Terrestrial vegetation of California*, 2nd Edition. Calif. Native Plant Soc. Spec. Publ. #9.

³⁷ Schoenherr, A. A. 1992. *A natural history of California*. University of California Press, Berkeley. 772p.

³⁸ Dale, N. 2000. Flowering plants of the Santa Monica Mountains. California Native Plant Society, 1722 J Street, Suite 17, Sacramento, CA 95814.

³⁹ Ballmer, G. R. 1995. What’s bugging coastal sage scrub. *Fremontia* 23(4):17-26.

⁴⁰ Root, R. B. 1967. The niche exploitation pattern of the blue-gray gnatcatcher. *Ecol. Monog.* 37:317-350.

the Santa Monica Mountains⁴¹. Five species of hummingbirds also follow the flowering cycle⁴².

Many species of 'opportunistic foragers', which utilize several different community types, perform important ecological roles during their seasonal movements. The scrub jay is a good example of such a species. The scrub jay is an omnivore and forages in coastal sage scrub, chaparral, and oak woodlands for insects, berries and notably acorns. Its foraging behavior includes the habit of burying acorns, usually at sites away from the parent tree canopy. Buried acorns have a much better chance of successful germination (about two-fold) than exposed acorns because they are protected from desiccation and predators. One scrub jay will bury approximately 5000 acorns in a year. The scrub jay therefore performs the function of greatly increasing recruitment and regeneration of oak woodland, a valuable and sensitive habitat type⁴³.

Like the scrub jay, most of the species of birds that inhabit the Mediterranean ecosystem in the Santa Monica Mountains require more than one community type in order to flourish. Many species include several community types in their daily activities. Other species tend to move from one community to another seasonally. The importance of maintaining the integrity of the multi-community ecosystem is clear in the following observations of Dr. Hartmut Walter of the University of California at Los Angeles:

"Bird diversity is directly related to the habitat mosaic and topographic diversity of the Santa Monicas. Most bird species in this bio-landscape require more than one habitat for survival and reproduction." "A significant proportion of the avifauna breeds in the wooded canyons of the Santa Monicas. Most of the canyon breeders forage every day in the brush- and grass-covered slopes, ridges and mesas. They would not breed in the canyons in the absence of the surrounding shrublands. Hawks, owls, falcons, orioles, flycatchers, woodpeckers, warblers, hummingbirds, etc. belong to this group. Conversely, some of the characteristic chaparral birds such as thrashers, quails, and wrentits need the canyons for access to shelter, protection from fire, and water. The regular and massive movement of birds between riparian corridors and adjacent shrublands has been demonstrated by qualitative and quantitative observations by several UCLA students⁴⁴."

Thus, the Mediterranean ecosystem of the Santa Monica Mountains is a mosaic of vegetation types linked together ecologically. The high biodiversity of the area results

⁴¹ Letter from Dr. Marti Witter, NPS, dated Sept. 13, 2001, in letters received and included in the September 2002 staff report for the Malibu LCP.

⁴² National Park Service. 1993. A checklist of the birds of the Santa Monica Mountains National Recreation Area. Southwest Parks and Monuments Assoc., 221 N. Court, Tucson, AZ. 85701

⁴³ Borchert, M. I., F. W. Davis, J. Michaelsen and L. D. Oyler. 1989. Interactions of factors affecting seedling recruitment of blue oak (*Quercus douglasii*) in California. Ecology 70:389-404. Bossema, I. 1979. Jays and oaks: An eco-ethological study of a symbiosis. Behavior 70:1-118. Schoenherr, A. A. 1992. A natural history of California. University of California Press, Berkeley. 772p.

⁴⁴ Walter, Hartmut. Bird use of Mediterranean habitats in the Santa Monica Mountains, Coastal Commission Workshop on the Significance of Native Habitats in the Santa Monica Mountains. CCC Hearing, June 13, 2002, Queen Mary Hotel.

from both the diversity and the interconnected nature of this mosaic. Most raptor species, for example, require large areas and will often require different habitats for perching, nesting and foraging. Fourteen species of raptors (13 of which are considered sensitive) are reported from the Santa Monica Mountains. These species utilize a variety of habitats including rock outcrops, oak woodlands, riparian areas, grasslands, chaparral, coastal sage scrub, estuaries and freshwater lakes⁴⁵.

When the community mosaic is disrupted and fragmented by development, many chaparral-associated native bird species are impacted. In a study of landscape-level fragmentation in the Santa Monica Mountains, Stralberg⁴⁶ found that the ash-throated flycatcher, Bewick's wren, wrentit, blue-gray gnatcatcher, California thrasher, orange-crowned warbler, rufous-crowned sparrow, spotted towhee, and California towhee all decreased in numbers as a result of urbanization. Soule⁴⁷ observed similar effects of fragmentation on chaparral and coastal sage scrub birds in the San Diego area.

In summary, all of the vegetation types in this ecosystem are strongly linked by animal movement and foraging. Whereas classification and mapping of vegetation types may suggest a snapshot view of the system, the seasonal movements and foraging of animals across these habitats illustrates the dynamic nature and vital connections that are crucial to the survival of this ecosystem.

Coastal Sage Scrub

"Coastal sage scrub" is a generic vegetation type that is inclusive of several subtypes⁴⁸. In the Santa Monica Mountains, coastal sage scrub is mostly of the type termed "Venturan Coastal Sage Scrub." In general, coastal sage scrub is comprised of dominant species that are semi-woody and low-growing, with shallow, dense roots that enable them to respond quickly to rainfall. Under the moist conditions of winter and spring, they grow quickly, flower, and produce light, wind-dispersed seeds, making them good colonizers following disturbance. These species cope with summer drought by dying back, dropping their leaves or producing a smaller summer leaf in order to reduce water loss. Stands of coastal sage scrub are much more open than chaparral and contain a greater admixture of herbaceous species. Coastal sage scrub is generally restricted to drier sites, such as low foothills, south-facing slopes, and shallow soils at higher elevations.

⁴⁵ National Park Service. 1993. A checklist of the birds of the Santa Monica Mountains National Recreation Area. Southwest Parks and Monuments Assoc., 221 N. Court, Tucson, AZ. 85701. and Letter from Dr. Marti Witter, NPS, Dated Sept. 13, 2001, in letters received and included in the September 2002 staff report for the Malibu LCP.

⁴⁶ Stralberg, D. 2000. Landscape-level urbanization effects on chaparral birds: A Santa Monica Mountains case study. p 125-136 in: Keeley, J. E., M. Baer-Keeley and C. J. Fotheringham (eds), 2nd Interface Between Ecology and Land Development in California, U.S. Geological Survey Open-File Report 00-62.

⁴⁷ Soule, M. E, D. T. Bolger, A. C. Alberts, J. Wright, M. Soric and S. Hill. 1988. Reconstructed dynamics of rapid extinctions of chaparral-requiring birds in urban habitat islands. *Conserv. Biol.* 2: 75-92.

⁴⁸ Kirkpatrick, J.B. and C.F. Hutchinson. 1977. The community composition of Californian coastal sage scrub. *Vegetatio* 35:21-33; Holland, 1986. op.cit.; Sawyer and Keeler-Wolf, 1995, op.cit.

The species composition and structure of individual stands of coastal sage scrub depend on moisture conditions that derive from slope, aspect, elevation and soil type. Drier sites are dominated by more drought-resistant species (e.g., California sagebrush, coast buckwheat, and *Opuntia* cactus). Where more moisture is available (e.g., north-facing slopes), larger evergreen species such as toyon, laurel sumac, lemonade berry, and sugar bush are common. As a result, there is more cover for wildlife, and movement of large animals from chaparral into coastal sage scrub is facilitated in these areas. Characteristic wildlife in this community includes Anna's hummingbirds, rufous-sided towhees, California quail, greater roadrunners, Bewick's wrens, coyotes, and coast horned lizards⁴⁹, but most of these species move between coastal sage scrub and chaparral during their daily activities or on a seasonal basis.

Of the many important ecosystem roles performed by the coastal sage scrub community, five are particularly important in the Santa Monica Mountains. Coastal sage scrub provides critical linkages between riparian corridors, provides essential habitat for species that require several habitat types during the course of their life histories, provides essential habitat for local endemics, supports rare species that are in danger of extinction, and reduces erosion, thereby protecting the water quality of coastal streams.

Riparian woodlands are primary contributors to the high biodiversity of the Santa Monica Mountains. The ecological integrity of those riparian habitats not only requires wildlife dispersal along the streams, but also depends on the ability of animals to move from one riparian area to another. Such movement requires that the riparian corridors be connected by suitable habitat. In the Santa Monica Mountains, coastal sage scrub and chaparral provide that function. Significant development in coastal sage scrub would reduce the riparian corridors to linear islands of habitat with severe edge effects⁵⁰, reduced diversity, and lower productivity.

Most wildlife species and many species of plants utilize several types of habitat. Many species of animals endemic to Mediterranean habitats move among several plant communities during their daily activities and many are reliant on different communities either seasonally or during different stages of their life cycle. Without an intact mosaic of coastal sage scrub, chaparral, and riparian community types, many species will not thrive. Specific examples of the importance of interconnected communities, or habitats, were provided in the discussion above. This is an essential ecosystem role of coastal sage scrub.

A characteristic of the coastal sage scrub vegetation type is a high degree of endemism. This is consonant with Westman's observation that 44 percent of the species he sampled in coastal sage scrub occurred at only one of his 67 sites, which were

⁴⁹ National Park Service. 2000. Draft: General Management Plan & Environmental Impact Statement, Santa Monica Mountains National Recreation Area, US Dept. of Interior, National Park Service, December 2000.

⁵⁰ Environmental impacts are particularly severe at the interface between development and natural habitats. The greater the amount of this "edge" relative to the area of natural habitat, the worse the impact.

distributed from the San Francisco Bay area to Mexico⁵¹. Species with restricted distributions are by nature more susceptible to loss or degradation of their habitat. Westman said of this unique and local aspect of coastal sage scrub species in California:

“While there are about 50 widespread sage scrub species, more than half of the 375 species encountered in the present study of the sage scrub flora are rare in occurrence within the habitat range. In view of the reduction of the area of coastal sage scrub in California to 10-15% of its former extent and the limited extent of preserves, measures to conserve the diversity of the flora are needed.”⁵²

Coastal sage scrub in southern California provides habitat for about 100 rare species⁵³, many of which are also endemic to limited geographic regions⁵⁴. In the Santa Monica Mountains, rare animals that inhabit coastal sage scrub⁵⁵ include the Santa Monica shieldback katydid, silvery legless lizard, coastal cactus wren, Bell's sparrow, San Diego desert woodrat, southern California rufous-crowned sparrow, coastal western whiptail, and San Diego horned lizard. Some of these species are also found in chaparral⁵⁶. Rare plants found in coastal sage scrub in the Santa Monica Mountains include Santa Susana tarplant, Coulter's saltbush, Blockman's dudleya, Braunton's milkvetch, Parry's spineflower, and Plummer's mariposa lily⁵⁷. A total of 32 sensitive species of reptiles, birds and mammals have been identified in this community by the National Park Service.⁵⁸

One of the most important ecological functions of coastal sage scrub in the Santa Monica Mountains is to protect water quality in coastal streams by reducing erosion in the watershed. Although shallow rooted, the shrubs that define coastal sage scrub have dense root masses that hold the surface soils much more effectively than the exotic annual grasses and forbs that tend to dominate in disturbed areas. The native shrubs of this community are resistant not only to drought, as discussed above, but well adapted to fire. Most of the semi-woody shrubs have some ability to crown sprout after

⁵¹ Westman, W.E. 1981. Diversity relations and succession in Californian coastal sage scrub. *Ecology* 62:170-184.

⁵² Ibid.

⁵³ Atwood, J. L. 1993. California gnatcatchers and coastal sage scrub: The biological basis for endangered species listing. pp.149-166 *In: Interface Between Ecology and Land Development in California*. Ed. J. E. Keeley, So. Calif. Acad. of Sci., Los Angeles. California Department of Fish and Game (CDFG). 1993. The Southern California Coastal Sage Scrub (CSS) Natural Communities Conservation Plan (NCCP). CDFG and Calif. Resources Agency, 1416 9th St., Sacramento, CA 95814.

⁵⁴ Westman, W.E. 1981. op. cit.

⁵⁵ Biological Resources Assessment of the Proposed Santa Monica Mountains Significant Ecological Area. Nov. 2000. Los Angeles Co., Dept. of Regional Planning, 320 West Temple St., Rm. 1383, Los Angeles, CA 90012.

⁵⁶ O'Leary J.F., S.A. DeSimone, D.D. Murphy, P.F. Brussard, M.S. Gilpin, and R.F. Noss. 1994. Bibliographies on coastal sage scrub and related malacophyllous shrublands of other Mediterranean-type climates. *California Wildlife Conservation Bulletin* 10:1-51.

⁵⁷ Biological Resources Assessment of the Proposed Santa Monica Mountains Significant Ecological Area. Nov. 2000. Los Angeles Co., Dept. of Regional Planning, 320 West Temple St., Rm. 1383, Los Angeles, CA 90012.

⁵⁸ NPS, 2000, op cit.

fire. Several CSS species (e.g., *Eriogonum cinereum*) in the Santa Monica Mountains and adjacent areas resprout vigorously and other species growing near the coast demonstrate this characteristic more strongly than do individuals of the same species growing at inland sites in Riverside County.⁵⁹ These shrub species also tend to recolonize rapidly from seed following fire. As a result they provide persistent cover that reduces erosion.

In addition to performing extremely important roles in the Mediterranean ecosystem, the coastal sage scrub community type has been drastically reduced in area by habitat loss to development. In the early 1980's it was estimated that 85 to 90 percent of the original extent of coastal sage scrub in California had already been destroyed.⁶⁰ Losses since that time have been significant and particularly severe in the coastal zone.

Therefore, because of its increasing rarity, its important role in the functioning of the Santa Monica Mountains Mediterranean ecosystem, and its extreme vulnerability to development, coastal sage scrub within the Santa Monica Mountains meets the definition of ESHA under the Coastal Act.

Chaparral

Another shrub community in the Santa Monica Mountain Mediterranean ecosystem is chaparral. Like "coastal sage scrub," this is a generic category of vegetation. Chaparral species have deep roots (10s of ft) and hard waxy leaves, adaptations to drought that increase water supply and decrease water loss at the leaf surface. Some chaparral species cope more effectively with drought conditions than do desert plants⁶¹. Chaparral plants vary from about one to four meters tall and form dense, intertwining stands with nearly 100 percent ground cover. As a result, there are few herbaceous species present in mature stands. Chaparral is well adapted to fire. Many species regenerate mainly by crown sprouting; others rely on seeds which are stimulated to germinate by the heat and ash from fires. Over 100 evergreen shrubs may be found in chaparral⁶². On average, chaparral is found in wetter habitats than coastal sage scrub, being more common at higher elevations and on north facing slopes.

The broad category "northern mixed chaparral" is the major type of chaparral shown in the National Park Service map of the Santa Monica Mountains. However, northern mixed chaparral can be variously dominated by chamise, scrub oak or one of several species of manzanita or by ceanothus. In addition, it commonly contains woody vines and large shrubs such as mountain mahogany, toyon, hollyleaf redberry, and sugarbush⁶³. The rare red shank chaparral plant community also occurs in the Santa Monica Mountains. Although included within the category "northern mixed chaparral" in

⁵⁹ Dr. John O'Leary, SDSU, personal communication to Dr. John Dixon, CCC, July 2, 2002

⁶⁰ Westman, W.E. 1981. op. cit.

⁶¹ Dr. Stephen Davis, Pepperdine University. Presentation at the CCC workshop on the significance of native habitats in the Santa Monica Mountains. June 13, 2002.

⁶² Keely, J.E. and S.C. Keeley. Chaparral. Pages 166-207 in M.G. Barbour and W.D. Billings, eds. North American Terrestrial Vegetation. New York, Cambridge University Press.

⁶³ Ibid.

the vegetation map, several types of ceanothus chaparral are reported in the Santa Monica Mountains. Ceanothus chaparral occurs on stable slopes and ridges, and may be dominated by bigpod ceanothus, buck brush ceanothus, hoaryleaf ceanothus, or greenbark ceanothus. In addition to ceanothus, other species that are usually present in varying amounts are chamise, black sage, holly-leaf redberry, sugarbush, and coast golden bush⁶⁴.

Several sensitive plant species that occur in the chaparral of the Santa Monica Mountains area are: Santa Susana tarplant, Lyon's pentachaeta, marcescent dudleya, Santa Monica Mountains dudleya, Braunton's milk vetch and salt spring checkerbloom⁶⁵. Several occurring or potentially occurring sensitive animal species in chaparral from the area are: Santa Monica shieldback katydid, western spadefoot toad, silvery legless lizard, San Bernardino ring-neck snake, San Diego mountain kingsnake, coast patch-nosed snake, sharp-shinned hawk, southern California rufous-crowned sparrow, Bell's sparrow, yellow warbler, pallid bat, long-legged myotis bat, western mastiff bat, and San Diego desert woodrat.⁶⁶

Coastal sage scrub and chaparral are the predominant generic community types of the Santa Monica Mountains and provide the living matrix within which rarer habitats like riparian woodlands exist. These two shrub communities share many important ecosystem roles. Like coastal sage scrub, chaparral within the Santa Monica Mountains provides critical linkages among riparian corridors, provides essential habitat for species that require several habitat types during the course of their life histories, provides essential habitat for sensitive species, and stabilizes steep slopes and reduces erosion, thereby protecting the water quality of coastal streams.

Many species of animals in Mediterranean habitats characteristically move among several plant communities during their daily activities, and many are reliant on different communities either seasonally or during different stages of their life cycle. The importance of an intact mosaic of coastal sage scrub, chaparral, and riparian community types is perhaps most critical for birds. However, the same principles apply to other taxonomic groups. For example, whereas coastal sage scrub supports a higher diversity of native ant species than chaparral, chaparral habitat is necessary for the coast horned lizard, an ant specialist⁶⁷. Additional examples of the importance of an interconnected communities, or habitats, were provided in the discussion of coastal sage scrub above. This is an extremely important ecosystem role of chaparral in the Santa Monica Mountains.

Chaparral is also remarkably adapted to control erosion, especially on steep slopes. The root systems of chaparral plants are very deep, extending far below the surface and

⁶⁴ Ibid.

⁶⁵ Biological Resources Assessment of the Proposed Santa Monica Mountains Significant Ecological Area. Nov. 2000. Los Angeles Co., Dept. of Regional Planning, 320 West Temple St., Rm. 1383, Los Angeles, CA 90012.

⁶⁶ Ibid.

⁶⁷ A.V. Suarez. Ants and lizards in coastal sage scrub and chaparral. A presentation at the CCC workshop on the significance of native habitats in the Santa Monica Mountains. June 13, 2002.

penetrating the bedrock below⁶⁸, so chaparral literally holds the hillsides together and prevents slippage.⁶⁹ In addition, the direct soil erosion from precipitation is also greatly reduced by 1) water interception on the leaves and above ground foliage and plant structures, and 2) slowing the runoff of water across the soil surface and providing greater soil infiltration. Chaparral plants are extremely resistant to drought, which enables them to persist on steep slopes even during long periods of adverse conditions. Many other species die under such conditions, leaving the slopes unprotected when rains return. Since chaparral plants recover rapidly from fire, they quickly re-exert their ground stabilizing influence following burns. The effectiveness of chaparral for erosion control after fire increases rapidly with time⁷⁰. Thus, the erosion from a 2-inch rain-day event drops from 5 yd³/acre of soil one year after a fire to 1 yd³/acre after 4 years.⁷¹ The following table illustrates the strong protective effect of chaparral in preventing erosion.

Soil erosion as a function of 24-hour precipitation and chaparral age.

Years Since Fire	Erosion (yd ³ /acre) at Maximum 24-hr Precipitation of:		
	2 inches	5 inches	11 inches
1	5	20	180
4	1	12	140
17	0	1	28
50+	0	0	3

Therefore, because of its important roles in the functioning of the Santa Monica Mountains Mediterranean ecosystem, and its extreme vulnerability to development, chaparral within the Santa Monica Mountains meets the definition of ESHA under the Coastal Act.

Oak Woodland and Savanna

Coast live oak woodland occurs mostly on north slopes, shaded ravines and canyon bottoms. Besides the coast live oak, this plant community includes hollyleaf cherry, California bay laurel, coffeeberry, and poison oak. Coast live oak woodland is more

⁶⁸ Helmers, H., J.S. Horton, G. Juhren and J. O'Keefe. 1955. Root systems of some chaparral plants in southern California. *Ecology* 36(4):667-678. Kummerow, J. and W. Jow. 1977. Root systems of chaparral shrubs. *Oecologia* 29:163-177.

⁶⁹ Radtke, K. 1983. *Living more safely in the chaparral-urban interface*. General Technical Report PSW-67. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Berkeley, California. 51 pp.

⁷⁰ Kittredge, J. 1973. *Forest influences — the effects of woody vegetation on climate, water, and soil*. Dover Publications, New York. 394 pp. Longcore, T and C. Rich. 2002. Protection of environmentally sensitive habitat areas in proposed local coastal plan for the Santa Monica Mountains. (Table 1). The Urban Wildlands Group, Inc., P.O. Box 24020 Los Angeles, CA 90024. Vicars, M. (ed.) 1999. *FireSmart: protecting your community from wildfire*. Partners in Protection, Edmonton, Alberta.

⁷¹ Ibid.

tolerant of salt-laden fog than other oaks and is generally found nearer the coast⁷². Coast live oak also occurs as a riparian corridor species within the Santa Monica Mountains.

Valley oaks are endemic to California and reach their southern most extent in the Santa Monica Mountains. Valley oaks were once widely distributed throughout California's perennial grasslands in central and coastal valleys. Individuals of this species may survive 400-600 years. Over the past 150 years, valley oak savanna habitat has been drastically reduced and altered due to agricultural and residential development. The understory is now dominated by annual grasses and recruitment of seedlings is generally poor. This is a very threatened habitat.

The important ecosystem functions of oak woodlands and savanna are widely recognized⁷³. These habitats support a high diversity of birds⁷⁴, and provide refuge for many species of sensitive bats⁷⁵. Typical wildlife in this habitat includes acorn woodpeckers, scrub jays, plain titmice, northern flickers, cooper's hawks, western screech owls, mule deer, gray foxes, ground squirrels, jackrabbits and several species of sensitive bats.

Therefore, because of their important ecosystem functions and vulnerability to development, oak woodlands and savanna within the Santa Monica Mountains met the definition of ESHA under the Coastal Act.

Grasslands

Grasslands consist of low herbaceous vegetation that is dominated by grass species but may also harbor native or non-native forbs.

California Perennial Grassland

Native grassland within the Santa Monica Mountains consists of perennial native needlegrasses: purple needlegrass, (*Nassella pulchra*), foothills needlegrass, (*Nassella lepida*) and nodding needlegrass (*Nassella cernua*). These grasses may occur in the same general area but they do not typically mix, tending to segregate based on slope

⁷² NPS 2000. op. cit.

⁷³ Block, W.M., M.L. Morrison, and J. Verner. 1990. Wildlife and oak-woodland interdependency. *Fremontia* 18(3):72-76. Pavlik, B.M., P.C. Muick, S. Johnson, and M. Popper. 1991. *Oaks of California*. Cachuma Press and California Oak Foundation, Los Olivos, California. 184 pp.

⁷⁴ Cody, M.L. 1977. Birds. Pp. 223-231 in Thrower, N.J.W., and D.E. Bradbury (eds.). *Chile-California Mediterranean scrub atlas*. US/IBP Synthesis Series 2. Dowden, Hutchinson & Ross, Stroudsburg, Pennsylvania. National Park Service. 1993. A checklist of the birds of the Santa Monica Mountains National Recreation Area. Southwest Parks and Monuments Assoc., 221 N. Court, Tucson, AZ. 85701

⁷⁵ Miner, K.L., and D.C. Stokes. 2000. Status, conservation issues, and research needs for bats in the south coast bioregion. Paper presented at *Planning for biodiversity: bringing research and management together*, February 29, California State University, Pomona, California.

and substrate factors⁷⁶. Mixed with these native needlegrasses are many non-native annual species that are characteristic of California annual grassland⁷⁷. Native perennial grasslands are now exceedingly rare⁷⁸. In California, native grasslands once covered nearly 20 percent of the land area, but today are reduced to less than 0.1 percent⁷⁹. The California Natural Diversity Database (CNDDDB) lists purple needlegrass habitat as a community needing priority monitoring and restoration. The CNDDDB considers grasslands with 10 percent or more cover by purple needlegrass to be significant, and recommends that these be protected as remnants of original California prairie. Patches of this sensitive habitat occur throughout the Santa Monica Mountains where they are intermingled with coastal sage scrub, chaparral and oak woodlands.

Many of the raptors that inhabit the Santa Monica Mountains make use of grasslands for foraging because they provide essential habitat for small mammals and other prey. Grasslands adjacent to woodlands are particularly attractive to these birds of prey since they simultaneously offer perching and foraging habitat. Particularly noteworthy in this regard are the white-tailed kite, northern harrier, sharp-shinned hawk, Cooper's hawk, red-shouldered hawk, red-tailed hawk, golden eagle, American kestrel, merlin, and prairie falcon⁸⁰.

Therefore, because of their extreme rarity, important ecosystem functions, and vulnerability to development, California native perennial grasslands within the Santa Monica Mountains meet the definition of ESHA under the Coastal Act.

California Annual Grassland

The term "California annual grassland" has been proposed to recognize the fact that non-native annual grasses should now be considered naturalized and a permanent feature of the California landscape and should be acknowledged as providing important ecological functions. These habitats support large populations of small mammals and provide essential foraging habitat for many species of birds of prey. California annual grassland generally consists of dominant invasive annual grasses that are primarily of Mediterranean origin. The dominant species in this community include common wild oats (*Avena fatua*), slender oat (*Avena barbata*), red brome (*Bromus madritensis* ssp. *Rubens*), ripgut brome, (*Bromus diandrus*), and herbs such as black mustard (*Brassica nigra*), wild radish (*Raphanus sativus*) and sweet fennel (*Foeniculum vulgare*). Annual grasslands are located in patches throughout the Santa Monica Mountains in previously disturbed areas, cattle pastures, valley bottoms and along roadsides. While many of

⁷⁶ Sawyer, J. O. and T. Keeler-Wolf. 1995. A manual of California vegetation. California Native Plant Society, 1722 J St., Suite 17, Sacramento, CA 95814.

⁷⁷ Biological Resources Assessment of the Proposed Santa Monica Mountains Significant Ecological Area. Nov. 2000. Los Angeles Co., Dept. of Regional Planning, 320 West Temple St., Rm. 1383, Los Angeles, CA 90012.

⁷⁸ Noss, R.F., E.T. LaRoe III and J.M. Scott. 1995. Endangered ecosystems of the United States: a preliminary assessment of loss and degradation. Biological Report 28. National Biological Service, U.S. Dept. of Interior.

⁷⁹ NPS 2000. op. cit.

⁸⁰ NPS 2000. op. cit.

these patches are dominated by invasive non-native species, it would be premature to say that they are never sensitive or do not harbor valuable annual native species. A large number of native forbs also may be present in these habitats⁸¹, and many native wildflowers occur primarily in annual grasslands. In addition, annual grasslands are primary foraging areas for many sensitive raptor species in the area.

Inspection of California annual grasslands should be done prior to any impacts to determine if any rare native species are present or if any rare wildlife rely on the habitat and to determine if the site meets the Coastal Act ESHA criteria.

Effects of Human Activities and Development on Habitats within the Santa Monica Mountains

The natural habitats of the Santa Monica Mountains are highly threatened by current development pressure, fragmentation and impacts from the surrounding megalopolis. The developed portions of the Santa Monica Mountains represents the extension of this urbanization into natural areas. About 54% of the undeveloped Santa Monica Mountains are in private ownership⁸², and computer simulation studies of the development patterns over the next 25 years predict a serious increase in habitat fragmentation⁸³. Development and associated human activities have many well-documented deleterious effects on natural communities. These environmental impacts may be both direct and indirect and include the effects of increased fire frequency, of fire clearance, of introduction of exotic species, and of night lighting.

Increased Fire Frequency

Since 1925, all the major fires in the Santa Monica Mountains have been caused by human activities⁸⁴. Increased fire frequency alters plant communities by creating conditions that select for some species over others. Strong resprouting plant species such as laurel sumac, are favored while non-sprouters like bigpod ceanothus, are at a disadvantage. Frequent fire recurrence before the non-sprouters can develop and reestablish a seed bank is detrimental, so that with each fire their chances for propagation are further reduced. Resprouters can be sending up new shoots quickly, and so they are favored in an increased fire frequency regime. Also favored are weedy and invasive species. Dr. Steven Davis in his abstract for a Coastal Commission

⁸¹ Holstein, G. 2001. Pre-agricultural grassland in Central California. *Madrono* 48(4):253-264. Stromberg, M.R., P. Kephart and V. Yadon. 2001. Composition, invasibility and diversity of coastal California grasslands. *Madrono* 48(4):236-252.

⁸² National Park Service. 2000. Draft: General Management Plan & Environmental Impact Statement, Santa Monica Mountains National Recreation Area, US Dept. of Interior, National Park Service, December 2000.

⁸³ Swenson, J. J., and J. Franklin. 2000. The effects of future urban development on habitat fragmentation in the Santa Monica Mountains. *Landscape Ecol.* 15:713-730.

⁸⁴ NPS, 2000, op. cit.

Workshop stated⁸⁵ *“We have evidence that recent increases in fire frequency has eliminated drought-hardy non-sprouters from chaparral communities near Malibu, facilitating the invasion of exotic grasses and forbs that further exacerbate fire frequency.”* Thus, simply increasing fire frequency from about once every 22 years (the historical frequency) to about once every 12 years (the current frequency) can completely change the vegetation community. This has cascading effects throughout the ecosystem.

Fuel Clearance

The removal of vegetation for fire protection in the Santa Monica Mountains is required by law in “Very High Fire Hazard Severity Zones”⁸⁶. Fuel removal is reinforced by insurance carriers⁸⁷. Generally, the Santa Monica Mountains are considered to be a high fire hazard severity zone. In such high fire hazard areas, homeowners must often resort to the California FAIR Plan to obtain insurance. Because of the high risk, all homes in “brush areas” are assessed an insurance surcharge if they have less than the recommended 200-foot fuel modification zone⁸⁸ around the home. The combination of insurance incentives and regulation assures that the 200-foot clearance zone will be applied universally⁸⁹. While it is not required that all of this zone be cleared of vegetation, the common practice is simply to disk this zone, essentially removing or highly modifying all native vegetation. For a new structure not adjacent to existing structures, this results in the removal or modification of a minimum of three acres of vegetation⁹⁰. While the directly impacted area is large, the effects of fuel modification extend beyond the 200-foot clearance area.

Effects of Fuel Clearance on Bird Communities

The impacts of fuel clearance on bird communities was studied by Stralberg who identified three ecological categories of birds in the Santa Monica Mountains: 1) local and long distance migrators (ash-throated flycatcher, Pacific-slope flycatcher, phainopepla, black-headed grosbeak), 2) chaparral-associated species (Bewick’s wren, wrentit, blue-gray gnatcatcher, California thrasher, orange-crowned warbler, rufous-crowned sparrow, spotted towhee, California towhee) and 3) urban-associated species

⁸⁵ Davis, Steven. Effects of fire and other factors on patterns of chaparral in the Santa Monica Mountains, Coastal Commission Workshop on the Significance of Native Habitats in the Santa Monica Mountains. CCC Hearing, June 13, 2002, Queen Mary Hotel.

⁸⁶ 1996 Los Angeles County Fire Code Section 1117.2.1

⁸⁷ Longcore, T and C. Rich. 2002. Protection of environmentally sensitive habitat areas in proposed local coastal plan for the Santa Monica Mountains. The Urban Wildlands Group, Inc., P.O. Box 24020 Los Angeles, CA 90024. Vicars, M. (ed.) 1999. FireSmart: protecting your community from wildfire. Partners in Protection, Edmonton, Alberta.

⁸⁸ Fuel Modification Plan Guidelines. Co. of Los Angeles Fire Department, Fuel Modification Unit, Prevention Bureau, Forestry Division, Brush Clearance Section, January 1998.

⁸⁹ Longcore, T and C. Rich. 2002. Protection of environmentally sensitive habitat areas in proposed local coastal plan for the Santa Monica Mountains. The Urban Wildlands Group, Inc., P.O. Box 24020 Los Angeles, CA 90024.

⁹⁰ Ibid.

(mourning dove, American crow, Western scrub-jay, Northern mockingbird)⁹¹. It was found in this study that the number of migrators and chaparral-associated species decreased due to habitat fragmentation while the abundance of urban-associated species increased. The impact of fuel clearance is to greatly increase this edge-effect of fragmentation by expanding the amount of cleared area and “edge” many-fold. Similar results of decreases in fragmentation-sensitive bird species are reported from the work of Bolger et al. in southern California chaparral⁹².

Effects of Fuel Clearance on Arthropod Communities

Fuel clearance and habitat modification may also disrupt native arthropod communities, and this can have surprising effects far beyond the cleared area on species seemingly unrelated to the direct impacts. A particularly interesting and well-documented example with ants and lizards illustrates this point. When non-native landscaping with intensive irrigation is introduced, the area becomes favorable for the invasive and non-native Argentine ant. This ant forms “super colonies” that can forage more than 650 feet out into the surrounding native chaparral or coastal sage scrub around the landscaped area⁹³. The Argentine ant competes with native harvester ants and carpenter ants displacing them from the habitat⁹⁴. These native ants are the primary food resource for the native coast horned lizard, a California “Species of Special Concern.” As a result of Argentine ant invasion, the coast horned lizard and its native ant food resources are diminished in areas near landscaped and irrigated developments⁹⁵. In addition to specific effects on the coast horned lizard, there are other Mediterranean habitat ecosystem processes that are impacted by Argentine ant invasion through impacts on long-evolved native ant-plant mutualisms⁹⁶. The composition of the whole arthropod community changes and biodiversity decreases when habitats are subjected to fuel modification. In coastal sage scrub disturbed by fuel modification, fewer arthropod

⁹¹ Stralberg, D. 2000. Landscape-level urbanization effects on chaparral birds: a Santa Monica Mountains case study. Pp. 125–136 in Keeley, J.E., M. Baer-Keeley, and C.J. Fotheringham (eds.). *2nd interface between ecology and land development in California*. U.S. Geological Survey, Sacramento, California.

⁹² Bolger, D. T., T. A. Scott and J. T. Rotenberry. 1997. Breeding bird abundance in an urbanizing landscape in coastal Southern California. *Conserv. Biol.* 11:406-421.

⁹³ Suarez, A.V., D.T. Bolger and T.J. Case. 1998. Effects of fragmentation and invasion on native ant communities in coastal southern California. *Ecology* 79(6):2041-2056.

⁹⁴ Holway, D.A. 1995. The distribution of the Argentine ant (*Linepithema humile*) in central California: a twenty-year record of invasion. *Conservation Biology* 9:1634-1637. Human, K.G. and D.M. Gordon. 1996. Exploitation and interference competition between the invasive Argentine ant, (*Linepithema humile*), and native ant species. *Oecologia* 105:405-412.

⁹⁵ Fisher, R.N., A.V. Suarez and T.J. Case. 2002. Spatial patterns in the abundance of the coastal horned lizard. *Conservation Biology* 16(1):205-215. Suarez, A.V. J.Q. Richmond and T.J. Case. 2000. Prey selection in horned lizards following the invasion of Argentine ants in southern California. *Ecological Applications* 10(3):711-725.

⁹⁶ Suarez, A.V., D.T. Bolger and T.J. Case. 1998. Effects of fragmentation and invasion on native ant communities in coastal southern California. *Ecology* 79(6):2041-2056. Bond, W. and P. Slingsby. Collapse of an Ant-Plant Mutualism: The Argentine Ant (*Iridomyrmex humilis*) and Myrmecochorous Proteaceae. *Ecology* 65(4):1031-1037.

predator species are seen and more exotic arthropod species are present than in undisturbed habitats⁹⁷.

Studies in the Mediterranean vegetation of South Africa (equivalent to California shrubland with similar plant species) have shown how the invasive Argentine ant can disrupt the whole ecosystem.⁹⁸ In South Africa the Argentine ant displaces native ants as they do in California. Because the native ants are no longer present to collect and bury seeds, the seeds of the native plants are exposed to predation, and consumed by seed eating insects, birds and mammals. When this habitat burns after Argentine ant invasion the large-seeded plants that were protected by the native ants all but disappear. So the invasion of a non-native ant species drives out native ants, and this can cause a dramatic change in the species composition of the plant community by disrupting long-established seed dispersal mutualisms. In California, some insect eggs are adapted to being buried by native ants in a manner similar to plant seeds⁹⁹.

Artificial Night Lighting

One of the more recently recognized human impacts on ecosystem function is that of artificial night lighting as it effects the behavior and function of many different types of organisms¹⁰⁰. For literally billions of years the only nighttime sources of light were the moon and stars, and living things have adapted to this previously immutable standard and often depend upon it for their survival. A review of lighting impacts suggests that whereas some species are unaffected by artificial night lighting, many others are severely impacted. Overall, most impacts are negative ones or ones whose outcome is unknown. Research to date has found negative impacts to plants, aquatic and terrestrial invertebrates, amphibians, fish, birds and mammals, and a detailed literature review can be found in the report by Longcore and Rich¹⁰¹.

Summary

In a past action, the Coastal Commission found¹⁰² that the Santa Monica Mountains Mediterranean Ecosystem, which includes the undeveloped native habitats of the Santa Monica Mountains, is rare and especially valuable because of its relatively pristine

⁹⁷ Longcore, T.R. 1999. Terrestrial arthropods as indicators of restoration success in coastal sage scrub. Ph.D. Dissertation, University of California, Los Angeles.

⁹⁸ Christian, C. 2001. Consequences of a biological invasion reveal the importance of mutualism for plant communities. *Nature* 413:635-639.

⁹⁹ Hughes, L. and M. Westoby. 1992. Capitula on stick insect eggs and elaiosomes on seeds: convergent adaptations for burial by ants. *Functional Ecology* 6:642-648.

¹⁰⁰ . Longcore, T and C. Rich. 2002. Protection of environmentally sensitive habitat areas in proposed local coastal plan for the Santa Monica Mountains. The Urban Wildlands Group, Inc., P.O. Box 24020 Los Angeles, CA 90024.

¹⁰¹ Ibid, and Ecological Consequences of Artificial Night Lighting, Conference, February 23-24, 2002, UCLA Los Angeles, California.

¹⁰² Revised Findings for the City of Malibu Local Coastal Program (as adopted on September 13, 2002) adopted on February 6, 2003.

character, physical complexity, and resultant biological diversity. The undeveloped native habitats within the Santa Monica Mountains that are discussed above are ESHA because of their valuable roles in that ecosystem, including providing a critical mosaic of habitats required by many species of birds, mammals and other groups of wildlife, providing the opportunity for unrestricted wildlife movement among habitats, supporting populations of rare species, and preventing the erosion of steep slopes and thereby protecting riparian corridors, streams and, ultimately, shallow marine waters.

The importance the native habitats in the Santa Monica Mountains was emphasized nearly 20 years ago by the California Department of Fish and Game¹⁰³. Commenting on a Draft Land Use Plan for the City of Malibu, the Regional Manager wrote that, "It is essential that large areas of land be reclassified to reflect their true status as ESHAs. One of the major needs of the Malibu LUP is that it should provide protection for entire drainages and not just stream bottoms." These conclusions were supported by the following observations:

"It is a fact that many of the wildlife species of the Santa Monica Mountains, such as mountain lion, deer, and raccoon, have established access routes through the mountains. They often travel to and from riparian zones and development such as high density residential may adversely affect a wildlife corridor.

Most animal species that exist in riparian areas will, as part of their life histories, also be found in other habitat types, including chapparal (sic) or grassland. For example, hawks nest and roost in riparian areas, but are dependent on large open areas for foraging. For the survival of many species, particularly those high on the food chain, survival will depend upon the presence of such areas. Such areas in the Santa Monica Mountains include grassland and coastal sage scrub communities, which have been documented in the SEA studies as supporting a wide diversity of plant and animal life."

This analysis by the Department of Fish and Game is consonant with the findings of the Commission in the case of the Malibu LCP, and with the conclusion that large contiguous areas of relatively pristine native habitat in the Santa Monica Mountains meet the definition of ESHA under the Coastal Act.

¹⁰³ Letter from F. A. Worthley, Jr. (CDFG) to N. Lucast (CCC) re Land Use Plan for Malibu dated March 22, 1983.